

## 32nd International Mathematical Olympiad

First Day — July 17, 1991

Time Limit:  $4\frac{1}{2}$  hours

1. Given a triangle  $ABC$ , let  $I$  be the center of its inscribed circle. The internal bisectors of the angles  $A, B, C$  meet the opposite sides in  $A', B', C'$  respectively. Prove that

$$\frac{1}{4} < \frac{AI \cdot BI \cdot CI}{AA' \cdot BB' \cdot CC'} \leq \frac{8}{27}.$$

2. Let  $n > 6$  be an integer and  $a_1, a_2, \dots, a_k$  be all the natural numbers less than  $n$  and relatively prime to  $n$ . If

$$a_2 - a_1 = a_3 - a_2 = \dots = a_k - a_{k-1} > 0,$$

prove that  $n$  must be either a prime number or a power of 2.

3. Let  $S = \{1, 2, 3, \dots, 280\}$ . Find the smallest integer  $n$  such that each  $n$ -element subset of  $S$  contains five numbers which are pairwise relatively prime.

Second Day — July 18, 1991

Time Limit:  $4\frac{1}{2}$  hours

1. Suppose  $G$  is a connected graph with  $k$  edges. Prove that it is possible to label the edges  $1, 2, \dots, k$  in such a way that at each vertex which belongs to two or more edges, the greatest common divisor of the integers labeling those edges is equal to 1.

[A *graph* consists of a set of points, called *vertices*, together with a set of *edges* joining certain pairs of distinct vertices. Each pair of vertices  $u, v$  belongs to at most one edge. The graph  $G$  is *connected* if for each pair of distinct vertices  $x, y$  there is some sequence of vertices  $x = v_0, v_1, v_2, \dots, v_m = y$  such that each pair  $v_i, v_{i+1}$  ( $0 \leq i < m$ ) is joined by an edge of  $G$ .]

2. Let  $ABC$  be a triangle and  $P$  an interior point of  $ABC$ . Show that at least one of the angles  $\angle PAB$ ,  $\angle PBC$ ,  $\angle PCA$  is less than or equal to  $30^\circ$ .

3. An infinite sequence  $x_0, x_1, x_2, \dots$  of real numbers is said to be *bounded* if there is a constant  $C$  such that  $|x_i| \leq C$  for every  $i \geq 0$ .

Given any real number  $a > 1$ , construct a bounded infinite sequence  $x_0, x_1, x_2, \dots$  such that

$$|x_i - x_j| |i - j|^a \geq 1$$

for every pair of distinct nonnegative integers  $i, j$ .