

Shortlist

2023

with solutions



IMO 2023



Chiba, JAPAN 64th

64th International Mathematical Olympiad
Chiba, Japan, 2nd–13th July 2023

64th International Mathematical Olympiad
Chiba, Japan, 2nd–13th July 2023



SHORTLISTED PROBLEMS

WITH SOLUTIONS

Note of Confidentiality

**The Shortlist has to be kept strictly confidential
until the conclusion of the following
International Mathematical Olympiad.**


IMO General Regulations §6.6

Contributing Countries

The Organising Committee and the Problem Selection Committee of IMO 2023 thank the following 52 countries for contributing 167 problem proposals:

Armenia, Australia, Austria, Azerbaijan, Bangladesh, Belarus, Belgium, Bulgaria, Brazil, Canada, China, Colombia, Croatia, Cyprus, Czech Republic, Estonia, Georgia, Germany, Greece, Hong Kong, Hungary, India, Iran, Israel, Latvia, Liechtenstein, Lithuania, Malaysia, Mexico, Mongolia, Morocco, Netherlands, New Zealand, North Macedonia, Poland, Portugal, Serbia, Singapore, Slovakia, Slovenia, South Africa, South Korea, Spain, Syria, Sweden, Taiwan, Tajikistan, Thailand, Turkey, Ukraine, United Kingdom, U.S.A., Uzbekistan.

Problem Selection Committee



Back: Yuya Matsumoto, Atsuo Yamauchi, Tetsushi Ito, Hiroki Kodama,
Ryu Minegishi, Genki Shimizu, Takuma Kitamura, Yuta Takaya, Dongryul Kim,
Front: Elisa Lorenzo García, Arnaud Maret, Paul Vaderlind, Tetsuya Ando,
Ivan Guo, Géza Kós, Sam Bealing

Problems

Algebra

A1. Professor Oak is feeding his 100 Pokémon. Each Pokémon has a bowl whose capacity is a positive real number of kilograms. These capacities are known to Professor Oak. The total capacity of all the bowls is 100 kilograms. Professor Oak distributes 100 kilograms of food in such a way that each Pokémon receives a non-negative integer number of kilograms of food (which may be larger than the capacity of their bowl). The *dissatisfaction level* of a Pokémon who received N kilograms of food and whose bowl has a capacity of C kilograms is equal to $|N - C|$.

Find the smallest real number D such that, regardless of the capacities of the bowls, Professor Oak can distribute the food in a way that the sum of the dissatisfaction levels over all the 100 Pokémon is at most D .

(Ukraine)

A2. Let \mathbb{R} be the set of real numbers. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that

$$f(x+y)f(x-y) \geq f(x)^2 - f(y)^2$$

for every $x, y \in \mathbb{R}$. Assume that the inequality is strict for some $x_0, y_0 \in \mathbb{R}$.

Prove that $f(x) \geq 0$ for every $x \in \mathbb{R}$ or $f(x) \leq 0$ for every $x \in \mathbb{R}$.

(Malaysia)

A3. Let $x_1, x_2, \dots, x_{2023}$ be *distinct* real positive numbers such that

$$a_n = \sqrt{(x_1 + x_2 + \dots + x_n) \left(\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n} \right)}$$

is an integer for every $n = 1, 2, \dots, 2023$. Prove that $a_{2023} \geq 3034$.

(Netherlands)

A4. Let $\mathbb{R}_{>0}$ be the set of positive real numbers. Determine all functions $f: \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ such that

$$x(f(x) + f(y)) \geq (f(f(x)) + y)f(y)$$

for every $x, y \in \mathbb{R}_{>0}$.

(Belgium)

A5. Let $a_1, a_2, \dots, a_{2023}$ be positive integers such that

- $a_1, a_2, \dots, a_{2023}$ is a permutation of $1, 2, \dots, 2023$, and
- $|a_1 - a_2|, |a_2 - a_3|, \dots, |a_{2022} - a_{2023}|$ is a permutation of $1, 2, \dots, 2022$.

Prove that $\max(a_1, a_{2023}) \geq 507$.

(Australia)

A6. Let $k \geq 2$ be an integer. Determine all sequences of positive integers a_1, a_2, \dots for which there exists a monic polynomial P of degree k with non-negative integer coefficients such that

$$P(a_n) = a_{n+1}a_{n+2} \cdots a_{n+k}$$

for every integer $n \geq 1$.

(Malaysia)

A7. Let N be a positive integer. Prove that there exist three permutations a_1, a_2, \dots, a_N ; b_1, b_2, \dots, b_N ; and c_1, c_2, \dots, c_N of $1, 2, \dots, N$ such that

$$\left| \sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} - 2\sqrt{N} \right| < 2023$$

for every $k = 1, 2, \dots, N$.

(China)

Combinatorics

C1. Let m and n be positive integers greater than 1. In each unit square of an $m \times n$ grid lies a coin with its tail-side up. A *move* consists of the following steps:

1. select a 2×2 square in the grid;
2. flip the coins in the top-left and bottom-right unit squares;
3. flip the coin in either the top-right or bottom-left unit square.

Determine all pairs (m, n) for which it is possible that every coin shows head-side up after a finite number of moves.

(Thailand)


C2. Determine the maximal length L of a sequence a_1, \dots, a_L of positive integers satisfying both the following properties:

- every term in the sequence is less than or equal to 2^{2023} , and
- there does not exist a consecutive subsequence a_i, a_{i+1}, \dots, a_j (where $1 \leq i \leq j \leq L$) with a choice of signs $s_i, s_{i+1}, \dots, s_j \in \{1, -1\}$ for which

$$s_i a_i + s_{i+1} a_{i+1} + \dots + s_j a_j = 0.$$

(Czech Republic)

C3. Let n be a positive integer. We arrange $1 + 2 + \dots + n$ circles in a triangle with n rows, such that the i^{th} row contains exactly i circles. The following figure shows the case $n = 6$.



In this triangle, a *ninja-path* is a sequence of circles obtained by repeatedly going from a circle to one of the two circles directly below it. In terms of n , find the largest value of k such that if one circle from every row is coloured red, we can always find a ninja-path in which at least k of the circles are red.

(Netherlands)

C4. Let $n \geq 2$ be a positive integer. Paul has a $1 \times n^2$ rectangular strip consisting of n^2 unit squares, where the i^{th} square is labelled with i for all $1 \leq i \leq n^2$. He wishes to cut the strip into several pieces, where each piece consists of a number of consecutive unit squares, and then *translate* (without rotating or flipping) the pieces to obtain an $n \times n$ square satisfying the following property: if the unit square in the i^{th} row and j^{th} column is labelled with a_{ij} , then $a_{ij} - (i + j - 1)$ is divisible by n .

Determine the smallest number of pieces Paul needs to make in order to accomplish this.

(U.S.A.)

C5. Elisa has 2023 treasure chests, all of which are unlocked and empty at first. Each day, Elisa adds a new gem to one of the unlocked chests of her choice, and afterwards, a fairy acts according to the following rules:


- if more than one chests are unlocked, it locks one of them, or
- if there is only one unlocked chest, it unlocks all the chests.

Given that this process goes on forever, prove that there is a constant C with the following property: Elisa can ensure that the difference between the numbers of gems in any two chests never exceeds C , regardless of how the fairy chooses the chests to lock.

(Israel)

C6. Let N be a positive integer, and consider an $N \times N$ grid. A *right-down path* is a sequence of grid cells such that each cell is either one cell to the right of or one cell below the previous cell in the sequence. A *right-up path* is a sequence of grid cells such that each cell is either one cell to the right of or one cell above the previous cell in the sequence.

Prove that the cells of the $N \times N$ grid cannot be partitioned into less than N right-down or right-up paths. For example, the following partition of the 5×5 grid uses 5 paths.



(Canada)

C7. The Imomi archipelago consists of $n \geq 2$ islands. Between each pair of distinct islands is a unique ferry line that runs in both directions, and each ferry line is operated by one of k companies. It is known that if any one of the k companies closes all its ferry lines, then it becomes impossible for a traveller, no matter where the traveller starts at, to visit all the islands exactly once (in particular, not returning to the island the traveller started at).

Determine the maximal possible value of k in terms of n .

(Ukraine)

Geometry

G1. Let $ABCDE$ be a convex pentagon such that $\angle ABC = \angle AED = 90^\circ$. Suppose that the midpoint of CD is the circumcentre of triangle ABE . Let O be the circumcentre of triangle ACD .

Prove that line AO passes through the midpoint of segment BE .

(Slovakia)

G2. Let ABC be a triangle with $AC > BC$. Let ω be the circumcircle of triangle ABC and let r be the radius of ω . Point P lies on segment AC such that $BC = CP$ and point S is the foot of the perpendicular from P to line AB . Let ray BP intersect ω again at D and let Q lie on line SP such that $PQ = r$ and S, P, Q lie on the line in that order. Finally, let the line perpendicular to CQ from A intersect the line perpendicular to DQ from B at E .

Prove that E lies on ω .

(Iran)

G3. Let $ABCD$ be a cyclic quadrilateral with $\angle BAD < \angle ADC$. Let M be the midpoint of the arc CD not containing A . Suppose there is a point P inside $ABCD$ such that $\angle ADB = \angle CPD$ and $\angle ADP = \angle PCB$.

Prove that lines AD, PM, BC are concurrent.

(Slovakia)

G4. Let ABC be an acute-angled triangle with $AB < AC$. Denote its circumcircle by Ω and denote the midpoint of arc CAB by S . Let the perpendicular from A to BC meet BS and Ω at D and $E \neq A$ respectively. Let the line through D parallel to BC meet line BE at L and denote the circumcircle of triangle BDL by ω . Let ω meet Ω again at $P \neq B$.

Prove that the line tangent to ω at P , and line BS intersect on the internal bisector of $\angle BAC$.

(Portugal)

G5. Let ABC be an acute-angled triangle with circumcircle ω and circumcentre O . Points $D \neq B$ and $E \neq C$ lie on ω such that $BD \perp AC$ and $CE \perp AB$. Let CO meet AB at X , and BO meet AC at Y .

Prove that the circumcircles of triangles BXD and CYE have an intersection on line AO .

(Malaysia)

G6. Let ABC be an acute-angled triangle with circumcircle ω . A circle Γ is internally tangent to ω at A and also tangent to BC at D . Let AB and AC intersect Γ at P and Q respectively. Let M and N be points on line BC such that B is the midpoint of DM and C is the midpoint of DN . Lines MP and NQ meet at K and intersect Γ again at I and J respectively. The ray KA meets the circumcircle of triangle IJK at $X \neq K$.

Prove that $\angle BXP = \angle CXQ$.

(United Kingdom)

G7. Let ABC be an acute, scalene triangle with orthocentre H . Let ℓ_a be the line through the reflection of B with respect to CH and the reflection of C with respect to BH . Lines ℓ_b and ℓ_c are defined similarly. Suppose lines ℓ_a , ℓ_b , and ℓ_c determine a triangle \mathcal{T} .

Prove that the orthocentre of \mathcal{T} , the circumcentre of \mathcal{T} and H are collinear.

(Ukraine)

G8. Let ABC be an equilateral triangle. Points A_1 , B_1 , C_1 lie inside triangle ABC such that triangle $A_1B_1C_1$ is scalene, $BA_1 = A_1C$, $CB_1 = B_1A$, $AC_1 = C_1B$ and

$$\angle BA_1C + \angle CB_1A + \angle AC_1B = 480^\circ.$$

Lines BC_1 and CB_1 intersect at A_2 ; lines CA_1 and AC_1 intersect at B_2 ; and lines AB_1 and BA_1 intersect at C_2 .

Prove that the circumcircles of triangles AA_1A_2 , BB_1B_2 , CC_1C_2 have two common points.

(U.S.A.)

Number Theory

N1. Determine all positive, composite integers n that satisfy the following property: if the positive divisors of n are $1 = d_1 < d_2 < \cdots < d_k = n$, then d_i divides $d_{i+1} + d_{i+2}$ for every $1 \leq i \leq k - 2$.

(Colombia)

N2. Determine all pairs (a, p) of positive integers with p prime such that $p^a + a^4$ is a perfect square.

(Bangladesh)

N3. For positive integers n and $k \geq 2$ define $E_k(n)$ as the greatest exponent r such that k^r divides $n!$. Prove that there are infinitely many n such that $E_{10}(n) > E_9(n)$ and infinitely many m such that $E_{10}(m) < E_9(m)$.

(Brazil)

N4. Let $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n$ be $2n$ positive integers such that the $n + 1$ products

$$\begin{aligned} & a_1 a_2 a_3 \cdots a_n, \\ & b_1 a_2 a_3 \cdots a_n, \\ & b_1 b_2 a_3 \cdots a_n, \\ & \vdots \\ & b_1 b_2 b_3 \cdots b_n \end{aligned}$$

form a strictly increasing arithmetic progression in that order. Determine the smallest positive integer that could be the common difference of such an arithmetic progression.

(Canada)

N5. Let $a_1 < a_2 < a_3 < \cdots$ be positive integers such that a_{k+1} divides $2(a_1 + a_2 + \cdots + a_k)$ for every $k \geq 1$. Suppose that for infinitely many primes p , there exists k such that p divides a_k . Prove that for every positive integer n , there exists k such that n divides a_k .

(Netherlands)

N6. A sequence of integers a_0, a_1, a_2, \dots is called *kawaii*, if $a_0 = 0, a_1 = 1$, and, for any positive integer n , we have

$$(a_{n+1} - 3a_n + 2a_{n-1})(a_{n+1} - 4a_n + 3a_{n-1}) = 0.$$

An integer is called *kawaii* if it belongs to a *kawaii* sequence.

Suppose that two consecutive positive integers m and $m + 1$ are both *kawaii* (not necessarily belonging to the same *kawaii* sequence). Prove that 3 divides m , and that $m/3$ is *kawaii*.

(China)

N7. Let a, b, c, d be positive integers satisfying

$$\frac{ab}{a+b} + \frac{cd}{c+d} = \frac{(a+b)(c+d)}{a+b+c+d}.$$

Determine all possible values of $a + b + c + d$.

(Netherlands)

N8. Let $\mathbb{Z}_{>0}$ be the set of positive integers. Determine all functions $f: \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ such that

$$f^{bf(a)}(a+1) = (a+1)f(b)$$

holds for all $a, b \in \mathbb{Z}_{>0}$, where $f^k(n) = f(f(\cdots f(n)\cdots))$ denotes the composition of f with itself k times.

(Taiwan)

Solutions

Algebra

A1. Professor Oak is feeding his 100 Pokémon. Each Pokémon has a bowl whose capacity is a positive real number of kilograms. These capacities are known to Professor Oak. The total capacity of all the bowls is 100 kilograms. Professor Oak distributes 100 kilograms of food in such a way that each Pokémon receives a non-negative integer number of kilograms of food (which may be larger than the capacity of their bowl). The *dissatisfaction level* of a Pokémon who received N kilograms of food and whose bowl has a capacity of C kilograms is equal to $|N - C|$.

Find the smallest real number D such that, regardless of the capacities of the bowls, Professor Oak can distribute the food in a way that the sum of the dissatisfaction levels over all the 100 Pokémon is at most D .

(Ukraine)

Answer: The answer is $D = 50$.

Solution 1. First, consider the situation where 99 bowls have a capacity of 0.5 kilograms and the last bowl has a capacity of 50.5 kilograms. No matter how Professor Oak distributes the food, the dissatisfaction level of every Pokémon will be at least 0.5. This amounts to a total dissatisfaction level of at least 50, proving that $D \geq 50$.

Now we prove that no matter what the capacities of the bowls are, Professor Oak can always distribute food in a way that the total dissatisfaction level is at most 50. We start by fixing some notation. We number the Pokémon from 1 to 100. Let $C_i > 0$ be the capacity of the bowl of the i^{th} Pokémon. By assumption, we have $C_1 + C_2 + \cdots + C_{100} = 100$. We write $F_i := C_i - \lfloor C_i \rfloor$ for the fractional part of C_i . Without loss of generality, we may assume that $F_1 \leq F_2 \leq \cdots \leq F_{100}$.

Here is a strategy: Professor Oak starts by giving $\lfloor C_i \rfloor$ kilograms of food to the i^{th} Pokémon. Let

$$R := 100 - \lfloor C_1 \rfloor - \lfloor C_2 \rfloor - \cdots - \lfloor C_{100} \rfloor = F_1 + F_2 + \cdots + F_{100} \geq 0$$

be the amount of food left. He continues by giving an extra kilogram of food to the R Pokémon numbered $100 - R + 1, 100 - R + 2, \dots, 100$, i.e. the Pokémon with the R largest values of F_i . By doing so, Professor Oak distributed 100 kilograms of food. The total dissatisfaction level with this strategy is

$$d := F_1 + \cdots + F_{100-R} + (1 - F_{100-R+1}) + \cdots + (1 - F_{100}).$$

We can rewrite

$$\begin{aligned} d &= 2(F_1 + \cdots + F_{100-R}) + R - (F_1 + \cdots + F_{100}) \\ &= 2(F_1 + \cdots + F_{100-R}). \end{aligned}$$

Now, observe that the arithmetic mean of $F_1, F_2, \dots, F_{100-R}$ is not greater than the arithmetic mean of F_1, F_2, \dots, F_{100} , because we assumed $F_1 \leq F_2 \leq \cdots \leq F_{100}$. Therefore

$$d \leq 2(100 - R) \cdot \frac{F_1 + \cdots + F_{100}}{100} = 2 \cdot \frac{R(100 - R)}{100}.$$

Finally, we use the AM-GM inequality to see that $R(100 - R) \leq \frac{100^2}{2^2}$ which implies $d \leq 50$. We conclude that there is always a distribution for which the total dissatisfaction level is at most 50, proving that $D \leq 50$.

Solution 2. We adopt the same notation as in Solution 1. Let $C_i > 0$ be the capacity of the bowl of the i^{th} Pokémon. By assumption, we have $C_1 + C_2 + \cdots + C_{100} = 100$. We write $F_i := C_i - \lfloor C_i \rfloor$ for the fractional part of C_i , and $R = F_1 + F_2 + \cdots + F_{100}$. Note that $R = 100 - \lfloor C_1 \rfloor - \cdots - \lfloor C_{100} \rfloor$ is an integer.

This solution uses the probabilistic method. We consider all distributions in which each Pokémon receives $\lfloor C_i \rfloor + \varepsilon_i$ kilograms of food, where $\varepsilon_i \in \{0, 1\}$ and $\varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_{100} = R$. There are $\binom{100}{R}$ such distributions. Suppose each of them occurs in an equal probability. In other words,

$$\varepsilon_i = \begin{cases} 0 & \text{with probability } \frac{100-R}{100}, \\ 1 & \text{with probability } \frac{R}{100}. \end{cases}$$

The expected value of the dissatisfaction level of the i^{th} Pokémon is

$$\frac{100-R}{100}(C_i - \lfloor C_i \rfloor) + \frac{R}{100}(\lfloor C_i \rfloor + 1 - C_i) = \frac{100-R}{100}F_i + \frac{R}{100}(1 - F_i).$$

Hence, the expected value of the total dissatisfaction level is

$$\begin{aligned} \sum_{i=1}^{100} \left(\frac{100-R}{100}F_i + \frac{R}{100}(1 - F_i) \right) &= \frac{100-R}{100} \sum_{i=1}^{100} F_i + \frac{R}{100} \sum_{i=1}^{100} (1 - F_i) \\ &= \frac{100-R}{100} \cdot R + \frac{R}{100} \cdot (100 - R) \\ &= 2 \cdot \frac{R(100-R)}{100}. \end{aligned}$$

As in Solution 1, this is at most 50. We conclude that there is at least one distribution for which the total dissatisfaction level is at most 50.

A2.

Let \mathbb{R} be the set of real numbers. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that

$$f(x+y)f(x-y) \geq f(x)^2 - f(y)^2$$

for every $x, y \in \mathbb{R}$. Assume that the inequality is strict for some $x_0, y_0 \in \mathbb{R}$.

Prove that $f(x) \geq 0$ for every $x \in \mathbb{R}$ or $f(x) \leq 0$ for every $x \in \mathbb{R}$.

(Malaysia)

Common remarks. We will say that f has constant sign, if f satisfies the conclusion of the problem.

Solution 1. We introduce the new variables $s := x + y$ and $t := x - y$. Equivalently, $x = \frac{s+t}{2}$ and $y = \frac{s-t}{2}$. The inequality becomes

$$f(s)f(t) \geq f\left(\frac{s+t}{2}\right)^2 - f\left(\frac{s-t}{2}\right)^2$$

for every $s, t \in \mathbb{R}$. We replace t by $-t$ to obtain

$$f(s)f(-t) \geq f\left(\frac{s-t}{2}\right)^2 - f\left(\frac{s+t}{2}\right)^2.$$

Summing the previous two inequalities gives

$$f(s)(f(t) + f(-t)) \geq 0$$

for every $s, t \in \mathbb{R}$. This inequality is strict for $s = x_0 + y_0$ and $t = x_0 - y_0$ by assumption. In particular, there exists some $t_0 = x_0 - y_0$ for which $f(t_0) + f(-t_0) \neq 0$. Since $f(s)(f(t_0) + f(-t_0)) \geq 0$ for every $s \in \mathbb{R}$, we conclude that $f(s)$ must have constant sign.

Solution 2. We do the same change of variables as in Solution 1 to obtain

$$f(s)f(t) \geq f\left(\frac{s+t}{2}\right)^2 - f\left(\frac{s-t}{2}\right)^2. \quad (1)$$

In this solution, we replace s by $-s$ (instead of t by $-t$). This gives

$$f(-s)f(t) \geq f\left(\frac{-s+t}{2}\right)^2 - f\left(\frac{-s-t}{2}\right)^2. \quad (2)$$

We now go back to the original inequality. Substituting $x = y$ gives $f(2x)f(0) \geq 0$ for every $x \in \mathbb{R}$. If $f(0) \neq 0$, then we conclude that f indeed has constant sign. From now on, we will assume that

$$f(0) = 0.$$

Substituting $x = -y$ gives $f(-x)^2 \geq f(x)^2$. By permuting x and $-x$, we conclude that

$$f(-x)^2 = f(x)^2$$

for every $x \in \mathbb{R}$.

Using the relation $f(x)^2 = f(-x)^2$, we can rewrite (2) as

$$f(-s)f(t) \geq f\left(\frac{s-t}{2}\right)^2 - f\left(\frac{s+t}{2}\right)^2.$$

Summing this inequality with (1), we obtain

$$(f(s) + f(-s))f(t) \geq 0$$

for every $s, t \in \mathbb{R}$ and we can conclude as in Solution 1.

Solution 3. We prove the contrapositive of the problem statement. Assume that there exist $a, b \in \mathbb{R}$ such that $f(a) < 0$ and $f(b) > 0$. We want to prove that the inequality is actually an equality, i.e. it is never strict.

Lemma 1. The function f is odd, i.e. $f(x) + f(-x) = 0$ for every $x \in \mathbb{R}$.

Proof. We plug in $x = \frac{a+u}{2}$ and $y = \frac{a-u}{2}$ in the original inequality, where u is a free variable. We obtain

$$f(a)f(u) \geq f\left(\frac{a+u}{2}\right)^2 - f\left(\frac{a-u}{2}\right)^2.$$

Replacing u with $-u$ and summing the two inequalities as in the previous solutions, we get

$$f(a)(f(u) + f(-u)) \geq 0$$

for every $u \in \mathbb{R}$. Since $f(a) < 0$ by assumption, we conclude that $f(u) + f(-u) \leq 0$ for every $u \in \mathbb{R}$.

We can repeat the above argument with b instead of a . Since $f(b) > 0$ by assumption, we conclude that $f(u) + f(-u) \geq 0$ for every $u \in \mathbb{R}$. This implies that $f(u) + f(-u) = 0$ for every $u \in \mathbb{R}$. \square

Now, using that f is odd, we can write the following chain of inequalities

$$\begin{aligned} f(x)^2 - f(y)^2 &\leq f(x+y)f(x-y) \\ &= -f(y+x)f(y-x) \\ &\leq -(f(y)^2 - f(x)^2) \\ &= f(x)^2 - f(y)^2. \end{aligned}$$

We conclude that every inequality above is actually an inequality, so

$$f(x+y)f(x-y) = f(x)^2 - f(y)^2$$

for every $x, y \in \mathbb{R}$.

Solution 4. As in Solution 3, we prove the contrapositive of the statement. Assume that there exist $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$. We want to prove that the inequality is actually an equality, i.e. it is never strict.

In this solution, we construct an argument by multiplying inequalities, rather than adding them as in Solutions 1-3.

Lemma 2. $f(b)f(-b) < 0$.

Proof. Let $x_1 := \frac{a+b}{2}$ and $y_1 := \frac{a-b}{2}$ so that $a = x_1 + y_1$ and $b = x_1 - y_1$. Plugging in $x = x_1$ and $y = y_1$, we obtain

$$0 > f(a)f(b) = f(x_1 + y_1)f(x_1 - y_1) \geq f(x_1)^2 - f(y_1)^2$$

which implies $f(x_1)^2 - f(y_1)^2 < 0$. Similarly, by plugging in $x = y_1$ and $y = x_1$, we get

$$f(a)f(-b) = f(y_1 + x_1)f(y_1 - x_1) \geq f(y_1)^2 - f(x_1)^2.$$

Using $f(x_1)^2 - f(y_1)^2 < 0$, we conclude $f(a)f(-b) > 0$. If we multiply the two inequalities $f(a)f(b) < 0$ and $f(a)f(-b) > 0$, we get $f(a)^2 f(b)f(-b) < 0$ and hence

$$f(b)f(-b) < 0.$$

Lemma 3. $f(x)f(-x) \leq 0$ for every $x \in \mathbb{R}$. \square

Proof. As in Solution 2, we prove that $f(x)^2 = f(-x)^2$ for every $x \in \mathbb{R}$ and we rewrite the original inequality as

$$f(s)f(t) \geq f\left(\frac{s+t}{2}\right)^2 - f\left(\frac{s-t}{2}\right)^2.$$

We replace s by $-s$ and t by $-t$, and use the relation $f(x)^2 = f(-x)^2$, to get

$$\begin{aligned} f(-s)f(-t) &\geq f\left(\frac{-s-t}{2}\right)^2 - f\left(\frac{-s+t}{2}\right)^2 \\ &= f\left(\frac{s+t}{2}\right)^2 - f\left(\frac{s-t}{2}\right)^2. \end{aligned}$$

Up to replacing t by $-t$, we can assume that $f\left(\frac{s+t}{2}\right)^2 - f\left(\frac{s-t}{2}\right)^2 \geq 0$. Multiplying the two previous inequalities leads to

$$f(s)f(-s)f(t)f(-t) \geq 0$$

for every $s, t \in \mathbb{R}$. This shows that $f(s)f(-s)$ (as a function of s) has constant sign. Since $f(b)f(-b) < 0$, we conclude that

$$f(x)f(-x) \leq 0$$

for every $x \in \mathbb{R}$. □

Lemma 3, combined with the relation $f(x)^2 = f(-x)^2$, implies $f(x) + f(-x) = 0$ for every $x \in \mathbb{R}$, i.e. f is odd. We conclude with the same argument as in Solution 3.

Comment. The presence of squares on the right-hand side of the inequality is not crucial as Solution 1 illustrates very well. However, it allows non-constant functions such as $f(x) = |x|$ to satisfy the conditions of the problem statement.

A3.

Let $x_1, x_2, \dots, x_{2023}$ be *distinct* real positive numbers such that

$$a_n = \sqrt{(x_1 + x_2 + \dots + x_n) \left(\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n} \right)}$$

is an integer for every $n = 1, 2, \dots, 2023$. Prove that $a_{2023} \geq 3034$.

(Netherlands)

Solution 1. We start with some basic observations. First note that the sequence $a_1, a_2, \dots, a_{2023}$ is increasing and thus, since all elements are integers, $a_{n+1} - a_n \geq 1$. We also observe that $a_1 = 1$ and

$$a_2 = \sqrt{(x_1 + x_2) \left(\frac{1}{x_1} + \frac{1}{x_2} \right)} > 2$$

by Cauchy-Schwarz inequality and using $x_1 \neq x_2$. So, $a_2 \geq 3$.

Now, we proceed to the main part of the argument. We observe that 3034 is about three halves of 2023. Motivated by this observation, we will prove the following.

Claim. If $a_{n+1} - a_n = 1$, then $a_{n+2} - a_{n+1} \geq 2$.

In other words, the sequence has to increase by at least 2 at least half of the times. Assuming the claim is true, since $a_1 = 1$, we would be done since

$$\begin{aligned} a_{2023} &= (a_{2023} - a_{2022}) + (a_{2022} - a_{2021}) + \dots + (a_2 - a_1) + a_1 \\ &\geq (2 + 1) \cdot 1011 + 1 \\ &= 3034. \end{aligned}$$

We now prove the claim. We start by observing that

$$\begin{aligned} a_{n+1}^2 &= (x_1 + \dots + x_{n+1}) \left(\frac{1}{x_1} + \dots + \frac{1}{x_{n+1}} \right) \\ &= (x_1 + \dots + x_n) \left(\frac{1}{x_1} + \dots + \frac{1}{x_n} \right) + 1 \\ &\quad + \frac{1}{x_{n+1}}(x_1 + \dots + x_n) + x_{n+1} \left(\frac{1}{x_1} + \dots + \frac{1}{x_n} \right) \\ &\geq a_n^2 + 1 + 2\sqrt{\frac{1}{x_{n+1}}(x_1 + \dots + x_n) \cdot x_{n+1} \left(\frac{1}{x_1} + \dots + \frac{1}{x_n} \right)} \\ &= a_n^2 + 1 + 2a_n \\ &= (a_n + 1)^2, \end{aligned}$$

where we used AM-GM to obtain the inequality. In particular, if $a_{n+1} = a_n + 1$, then

$$\frac{1}{x_{n+1}}(x_1 + \dots + x_n) = x_{n+1} \left(\frac{1}{x_1} + \dots + \frac{1}{x_n} \right). \quad (1)$$

Now, assume for the sake of contradiction that both $a_{n+1} = a_n + 1$ and $a_{n+2} = a_{n+1} + 1$ hold. In this case, (1) gives

$$\frac{1}{x_{n+2}}(x_1 + \dots + x_{n+1}) = x_{n+2} \left(\frac{1}{x_1} + \dots + \frac{1}{x_{n+1}} \right).$$

We can rewrite this relation as

$$\frac{x_{n+1}}{x_{n+2}} \left(\frac{1}{x_{n+1}}(x_1 + \dots + x_n) + 1 \right) = \frac{x_{n+2}}{x_{n+1}} \left(x_{n+1} \left(\frac{1}{x_1} + \dots + \frac{1}{x_n} \right) + 1 \right).$$

From (1) again, we conclude that $x_{n+1} = x_{n+2}$ which is a contradiction.

Solution 2. The trick is to compare a_{n+2} and a_n . Observe that

$$\begin{aligned}
 a_{n+2}^2 &= (x_1 + \cdots + x_{n+2}) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_{n+2}} \right) \\
 &= (x_1 + \cdots + x_n) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} \right) + (x_{n+1} + x_{n+2}) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right) \\
 &\quad + (x_1 + \cdots + x_n) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right) + (x_{n+1} + x_{n+2}) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} \right) \\
 &\geq a_n^2 + (x_{n+1} + x_{n+2}) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right) \\
 &\quad + 2\sqrt{(x_{n+1} + x_{n+2}) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right) (x_1 + \cdots + x_n) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} \right)} \\
 &= a_n^2 + (x_{n+1} + x_{n+2}) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right) + 2a_n \sqrt{(x_{n+1} + x_{n+2}) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right)},
 \end{aligned}$$

where we used AM-GM to obtain the inequality. Furthermore, we have

$$(x_{n+1} + x_{n+2}) \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right) > 4$$

because $x_{n+1} \neq x_{n+2}$ by assumption. Therefore, it follows that

$$a_{n+2}^2 > a_n^2 + 4 + 4a_n = (a_n + 2)^2.$$

Because a_{n+2} and a_n are both positive integers, we conclude that

$$a_{n+2} \geq a_n + 3.$$

A simple induction gives $a_{2k+1} \geq 3k + a_1$ for every $k \geq 0$. Since $a_1 = 1$, it follows that $a_{2k+1} \geq 3k + 1$. We get the desired conclusion for $k = 1011$. \square

Comment 1. A similar argument as in Solution 2 shows that $a_2 \geq 3$ and $a_{2k} \geq 3k$ for every $k \geq 1$. Actually, these lower bounds on a_n are sharp (at least for $n \leq 2023$). In other words, there exists a sequence of distinct values $x_1, \dots, x_{2023} > 0$ for which

$$a_n = \begin{cases} \frac{3n-1}{2} & \text{if } n \text{ is odd,} \\ \frac{3n}{2} & \text{if } n \text{ is even,} \end{cases}$$

for $n = 1, \dots, 2023$. The value of x_1 can be chosen arbitrarily. The next values can be obtained inductively by solving the quadratic equation

$$a_{n+1}^2 = a_n^2 + 1 + \left(\sum_{i=1}^n x_i \right) \frac{1}{x_{n+1}} + \left(\sum_{i=1}^n \frac{1}{x_i} \right) x_{n+1}$$

for x_{n+1} . Computation gives, for $n \geq 1$,

$$x_{n+1} = \begin{cases} \frac{3n}{2\left(\sum_{i=1}^n \frac{1}{x_i}\right)} & \text{if } n \text{ is even,} \\ \frac{6n+1 \pm 3\sqrt{n(3n+2)}}{2\left(\sum_{i=1}^n \frac{1}{x_i}\right)} & \text{if } n \text{ is odd.} \end{cases}$$

One can check (with the help of a computer), that the values x_1, \dots, x_{2023} obtained by choosing $x_1 = 1$ and “+” every time in the odd case are indeed distinct.

It is interesting to note that the discriminant always vanishes in the even case. This is a consequence of $a_{n+1} = a_n + 1$ being achieved as an equality case of AM-GM. Another cute observation is that the ratio x_2/x_1 is equal to the fourth power of the golden ratio.

Comment 2. The estimations in Solutions 1 and 2 can be made more efficiently if one applies the following form of the Cauchy–Schwarz inequality instead:

$$\sqrt{(a+b)(c+d)} \geq \sqrt{ac} + \sqrt{bd} \quad (2)$$

for arbitrary nonnegative numbers a, b, c, d . Equality occurs if and only if $a : c = b : d = (a+b) : (c+d)$.

For instance, by applying (2) to $a = x_1 + \cdots + x_n$, $b = x_{n+1}$, $c = \frac{1}{x_1} + \cdots + \frac{1}{x_n}$ and $d = \frac{1}{x_{n+1}}$ we get

$$\begin{aligned} a_{n+1} &= \sqrt{(x_1 + \cdots + x_n + x_{n+1}) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} + \frac{1}{x_{n+1}} \right)} \\ &\geq \sqrt{(x_1 + \cdots + x_n) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} \right)} + \sqrt{x_{n+1} \cdot \frac{1}{x_{n+1}}} = a_n + 1. \end{aligned}$$

A study of equality cases show that equality cannot occur twice in a row, as in Solution 1. Suppose that $a_{n+1} = a_n + 1$ and $a_{n+2} = a_{n+1} + 1$ for some index n . By the equality case in (2) we have

$$\frac{(x_1 + \cdots + x_n) + x_{n+1}}{\left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} \right) + \frac{1}{x_{n+1}}} = \frac{x_{n+1}}{1/x_{n+1}} = x_{n+1}^2 \quad \text{because } a_{n+1} = a_n + 1,$$

and

$$\frac{x_1 + \cdots + x_n + x_{n+1}}{\frac{1}{x_1} + \cdots + \frac{1}{x_n} + \frac{1}{x_{n+1}}} = \frac{x_{n+2}}{1/x_{n+2}} = x_{n+2}^2 \quad \text{because } a_{n+2} = a_{n+1} + 1.$$

The left-hand sides are the same, so $x_{n+1} = x_{n+2}$, but this violates the condition that x_{n+1} and x_{n+2} are distinct.

The same trick applies to Solution 2. We can compare a_n and a_{n+2} directly as

$$\begin{aligned} a_{n+2} &= \sqrt{(x_1 + \cdots + x_n + x_{n+1} + x_{n+2}) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} + \frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right)} \\ &\geq \sqrt{(x_1 + \cdots + x_n) \left(\frac{1}{x_1} + \cdots + \frac{1}{x_n} \right)} + \sqrt{(x_{n+1} + x_{n+2}) \cdot \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right)} \\ &= a_n + \sqrt{(x_{n+1} + x_{n+2}) \cdot \left(\frac{1}{x_{n+1}} + \frac{1}{x_{n+2}} \right)} \\ &\geq a_n + 2. \end{aligned}$$

In the last estimate, equality is not possible because x_{n+1} and x_{n+2} are distinct, so $a_{n+2} > a_n + 2$ and therefore $a_{n+2} \geq a_n + 3$.

A4. Let $\mathbb{R}_{>0}$ be the set of positive real numbers. Determine all functions $f: \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ such that

$$x(f(x) + f(y)) \geq (f(f(x)) + y)f(y)$$

for every $x, y \in \mathbb{R}_{>0}$.

(Belgium)

Answer: All functions $f(x) = \frac{c}{x}$ for some $c > 0$.

Solution 1. Let $f: \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ be a function that satisfies the inequality of the problem statement. We will write $f^k(x) = f(f(\cdots f(x)\cdots))$ for the composition of f with itself k times, with the convention that $f^0(x) = x$. Substituting $y = x$ gives

$$x \geq f^2(x).$$

Substituting $x = f(y)$ instead leads to $f(y) + f^2(y) \geq y + f^3(y)$, or equivalently

$$f(y) - f^3(y) \geq y - f^2(y).$$

We can generalise this inequality. If we replace y by $f^{n-1}(y)$ in the above inequality, we get

$$f^n(y) - f^{n+2}(y) \geq f^{n-1}(y) - f^{n+1}(y),$$

for every $y \in \mathbb{R}_{>0}$ and for every integer $n \geq 1$. In particular, $f^n(y) - f^{n+2}(y) \geq y - f^2(y) \geq 0$ for every $n \geq 1$. Hereafter consider even integers $n = 2m$. Observe that

$$y - f^{2m}(y) = \sum_{i=0}^{m-1} (f^{2i}(y) - f^{2i+2}(y)) \geq m(y - f^2(y)).$$

Since f takes positive values, it holds that $y - f^{2m}(y) < y$ for every $m \geq 1$. So, we have proved that $y > m(y - f^2(y))$ for every $y \in \mathbb{R}_{>0}$ and every $m \geq 1$. Since $y - f^2(y) \geq 0$, this holds if only if

$$f^2(y) = y$$

for every $y \in \mathbb{R}_{>0}$. The original inequality becomes

$$xf(x) \geq yf(y)$$

for every $x, y \in \mathbb{R}_{>0}$. Hence, $xf(x)$ is constant. We conclude that $f(x) = c/x$ for some $c > 0$.

We now check that all the functions of the form $f(x) = c/x$ are indeed solutions of the original problem. First, note that all these functions satisfy $f(f(x)) = c/(c/x) = x$. So it's sufficient to check that $xf(x) \geq yf(y)$, which is true since $c \geq c$.

Solution 2. Let $f: \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ be a function that satisfies the inequality of the problem statement. As in Solution 1, we prove that

$$f^n(y) \geq f^{n+2}(y)$$

for every $y \in \mathbb{R}_{>0}$ and every $n \geq 0$. Since f takes positive values, this implies that

$$yf(y) \geq f(y)f^2(y) \geq f^2(y)f^3(y) \geq \cdots$$

In other words, $yf(y) \geq f^n(y)f^{n+1}(y)$ for every $y \in \mathbb{R}_{>0}$ and every $n \geq 1$.

We replace x by $f^n(x)$ in the original inequality and get

$$f^n(x) - f^{n+2}(x) \geq \frac{yf(y) - f^n(x)f^{n+1}(x)}{f(y)}.$$

Using that $xf(x) \geq f^n(x)f^{n+1}(x)$, we obtain

$$f^n(x) - f^{n+2}(x) \geq \frac{yf(y) - xf(x)}{f(y)}$$

for every $n \geq 0$. The same trick as in Solution 1 gives

$$x > x - f^{2m}(x) = \sum_{i=0}^{m-1} (f^{2i}(x) - f^{2i+2}(x)) \geq m \cdot \frac{yf(y) - xf(x)}{f(y)}$$

for every $x, y \in \mathbb{R}_{>0}$ and every $m \geq 1$. Possibly permuting x and y , we may assume that $yf(y) - xf(x) \geq 0$ then the above inequality implies $xf(x) = yf(y)$. We conclude as in Solution 1.

A5.

Let $a_1, a_2, \dots, a_{2023}$ be positive integers such that

- $a_1, a_2, \dots, a_{2023}$ is a permutation of $1, 2, \dots, 2023$, and
- $|a_1 - a_2|, |a_2 - a_3|, \dots, |a_{2022} - a_{2023}|$ is a permutation of $1, 2, \dots, 2022$.

Prove that $\max(a_1, a_{2023}) \geq 507$.

(Australia)

Solution. For the sake of clarity, we consider and prove the following generalisation of the original problem (which is the case $N = 1012$):

Let N be a positive integer and $a_1, a_2, \dots, a_{2N-1}$ be positive integers such that

- $a_1, a_2, \dots, a_{2N-1}$ is a permutation of $1, 2, \dots, 2N - 1$, and
- $|a_1 - a_2|, |a_2 - a_3|, \dots, |a_{2N-2} - a_{2N-1}|$ is a permutation of $1, 2, \dots, 2N - 2$.

Then $a_1 + a_{2N-1} \geq N + 1$ and hence $\max(a_1, a_{2N-1}) \geq \lceil \frac{N+1}{2} \rceil$.

Now we proceed to the proof of the generalised statement. We introduce the notion of *score* of a number $a \in \{1, 2, \dots, 2N - 1\}$. The score of a is defined to be

$$s(a) := |a - N|.$$

Note that, by the triangle inequality,

$$|a - b| \leq |a - N| + |N - b| = s(a) + s(b).$$

Considering the sum $|a_1 - a_2| + |a_2 - a_3| + \dots + |a_{2N-2} - a_{2N-1}|$, we find that

$$\begin{aligned} (N - 1)(2N - 1) &= |a_1 - a_2| + |a_2 - a_3| + \dots + |a_{2N-2} - a_{2N-1}| \\ &\leq 2(s(a_1) + s(a_2) + \dots + s(a_{2N-1})) - (s(a_1) + s(a_{2N-1})) \\ &= 2N(N - 1) - (s(a_1) + s(a_{2N-1})). \end{aligned}$$

For the last equality we used that the numbers $s(a_1), s(a_2), \dots, s(a_{2N-1})$ are a permutation of $0, 1, 1, 2, 2, \dots, N - 1, N - 1$.

Hence, $s(a_1) + s(a_{2N-1}) \leq 2N(N - 1) - (N - 1)(2N - 1) = N - 1$. We conclude that

$$(N - a_1) + (N - a_{2N-1}) \leq s(a_1) + s(a_{2N-1}) \leq N - 1,$$

which implies $a_1 + a_{2N-1} \geq N + 1$.

Comment 1. In the case $N = 1012$, such a sequence with $\max(a_1, a_{2023}) = 507$ indeed exists:

$$507, 1517, 508, 1516, \dots, 1011, 1013, 1012, 2023, 1, 2022, 2, \dots, 1518, 506.$$

For a general even number N , a sequence with $\max(a_1, a_{2N-1}) = \lceil \frac{N+1}{2} \rceil$ can be obtained similarly. If $N \geq 3$ is odd, the inequality is not sharp, because $\max(a_1, a_{2N-1}) = \frac{N+1}{2}$ and $a_1 + a_{2N-1} \geq N + 1$ together imply $a_1 = a_{2N-1} = \frac{N+1}{2}$, a contradiction.

Comment 2. The formulation of the author’s submission was slightly different:

Author’s formulation. Consider a sequence of positive integers a_1, a_2, a_3, \dots such that the following conditions hold for all positive integers m and n :

- $a_{n+2023} = a_n + 2023$,
- If $|a_{n+1} - a_n| = |a_{m+1} - a_m|$, then $2023 \mid (n - m)$, and
- The sequence contains every positive integer.

Prove that $a_1 \geq 507$.

The two formulations are equivalent up to relatively trivial arguments. Suppose (a_n) is a sequence satisfying the author’s formulation. From the first and third conditions, we see that a_1, \dots, a_{2023} is a permutation of $1, \dots, 2023$. Moreover, the sequence $|a_i - a_{i+1}|$ for $i = 1, 2, \dots, 2022$ consists of positive integers ≤ 2022 and has pairwise distinct elements by the second condition. Hence, it is a permutation of $1, \dots, 2022$. It also holds that $a_1 > a_{2023}$, since if $a_1 < a_{2023}$ then $|a_{2024} - a_{2023}| = |2023 + a_1 - a_{2023}| \leq 2022$, which should be equal to $|a_i - a_{i+1}|$ for some $1 \leq i \leq 2022$, contradicting the second condition. This reduces the problem to the Shortlist formulation.

Conversely, if the numbers a_1, \dots, a_{2023} satisfy the conditions of the Shortlist formulation, then, after possibly reversing the sequence to ensure $a_1 > a_{2023}$, the sequence can be extended to an infinite sequence satisfying the conditions of the author’s formulation.

A6. Let $k \geq 2$ be an integer. Determine all sequences of positive integers a_1, a_2, \dots for which there exists a monic polynomial P of degree k with non-negative integer coefficients such that

$$P(a_n) = a_{n+1}a_{n+2} \cdots a_{n+k}$$

for every integer $n \geq 1$.

(Malaysia)

Answer: The sequence (a_n) must be an arithmetic progression consisting of positive integers with common difference $d \geq 0$, and $P(x) = (x + d) \cdots (x + kd)$.

Common remarks. The following arguments and observations are implicit in the solutions given below.

Suppose the sequence (a_n) is an arithmetic progression with common difference $d \geq 0$. Then it satisfies the condition with

$$P(x) = (x + d) \cdots (x + kd).$$

This settles one direction. Now suppose (a_n) is a sequence satisfying the condition. We will show that it is a non-decreasing arithmetic progression.

Since $P(x)$ has non-negative integer coefficients, it is strictly increasing on the positive real line. In particular, it holds that, for any positive integer x, y ,

$$P(x) < P(y) \iff x < y.$$

Furthermore, if the sequence (a_n) is eventually constant, then $P(x) = x^k$ and the sequence (a_n) is actually constant. Indeed, if $P(x)$ were not the polynomial x^k , then $P(a_n) = a_{n+1} \cdots a_{n+k}$ cannot be satisfied for n such that $a_n = \cdots = a_{n+k}$. By a descending induction, we conclude that (a_n) is constant. Thus we can restrict to the case (a_n) is not eventually constant.

Solution 1. We assume that (a_n) is not eventually constant.

Step 1. The first goal is to show that the sequence must be *increasing*, i.e. $a_n < a_{n+1}$ for all $n \geq 1$.

First, by comparing the two equalities

$$\begin{aligned} P(a_n) &= a_{n+1}a_{n+2} \cdots a_{n+k}, \\ P(a_{n+1}) &= a_{n+2} \cdots a_{n+k}a_{n+k+1}, \end{aligned}$$

we observe that

$$a_n < a_{n+1} \iff P(a_n) < P(a_{n+1}) \iff a_{n+1} < a_{n+k+1}, \quad (1)$$

$$a_n > a_{n+1} \iff P(a_n) > P(a_{n+1}) \iff a_{n+1} > a_{n+k+1}, \quad (2)$$

$$a_n = a_{n+1} \iff P(a_n) = P(a_{n+1}) \iff a_{n+1} = a_{n+k+1}. \quad (3)$$

Claim 1. $a_n \leq a_{n+1}$ for all $n \geq 1$.

Proof. Suppose, to the contrary, that $a_{n(0)-1} > a_{n(0)}$ for some $n(0) \geq 2$. We will give an infinite sequence of positive integers $n(0) < n(1) < \cdots$ satisfying

$$a_{n(i)-1} > a_{n(i)} \text{ and } a_{n(i)} > a_{n(i+1)}.$$

Then $a_{n(0)}, a_{n(1)}, a_{n(2)}, \dots$ is an infinite decreasing sequence of positive integers, which is absurd.

We construct such a sequence inductively. If we have chosen $n(i)$, then we let $n(i+1)$ be the smallest index larger than $n(i)$ such that $a_{n(i)} > a_{n(i+1)}$. Note that such an index always exists and satisfies $n(i) + 1 \leq n(i+1) \leq n(i) + k$ because $a_{n(i)} > a_{n(i)+k}$ by (2). We need to check that $a_{n(i+1)-1} > a_{n(i+1)}$. This is immediate if $n(i+1) = n(i) + 1$ by construction. If $n(i+1) \geq n(i) + 2$, then $a_{n(i+1)-1} \geq a_{n(i)}$ by minimality of $n(i+1)$, and so $a_{n(i+1)-1} \geq a_{n(i)} > a_{n(i+1)}$. \square

We are now ready to prove that the sequence a_n is increasing. Suppose $a_n = a_{n+1}$ for some $n \geq 1$. Then we also have $a_{n+1} = a_{n+k+1}$ by (3), and since the sequence is non-decreasing we have $a_n = a_{n+1} = a_{n+2} = \cdots = a_{n+k+1}$. We repeat the argument for $a_{n+k} = a_{n+k+1}$ and get that the sequence is eventually constant, which contradicts our assumption. Hence

$$a_n < a_{n+1} \text{ for all } n \geq 1.$$

Step 2. The next and final goal is to prove that the sequence a_n is an *arithmetic progression*. Observe that we can make differences of terms appear as follows

$$\begin{aligned} P(a_n) &= a_{n+1}a_{n+2} \cdots a_{n+k} \\ &= (a_n + (a_{n+1} - a_n))(a_n + (a_{n+2} - a_n)) \cdots (a_n + (a_{n+k} - a_n)). \end{aligned}$$

We will prove that, for n large enough, the sum

$$(a_{n+1} - a_n) + (a_{n+2} - a_n) + \cdots + (a_{n+k} - a_n)$$

is equal to the coefficient b of the term x^{k-1} in P . The argument is based on the following claim.

Claim 2. There exists a bound A with the following properties:

1. If (c_1, \dots, c_k) is a k -tuple of positive integers with $c_1 + \cdots + c_k > b$, then for every $x \geq A$ we have $P(x) < (x + c_1)(x + c_2) \cdots (x + c_k)$.
2. If (c_1, \dots, c_k) is a k -tuple of positive integers with $c_1 + \cdots + c_k < b$, then for every $x \geq A$ we have $P(x) > (x + c_1)(x + c_2) \cdots (x + c_k)$.

Proof. It suffices to show parts 1 and 2 separately, because then we can take the maximum of two bounds.

We first show part 1. For each single (c_1, \dots, c_k) such a bound A exists since

$$P(x) - (x + c_1)(x + c_2) \cdots (x + c_k) = (b - (c_1 + \cdots + c_k))x^{k-1} + (\text{terms of degree } \leq k-2)$$

has negative leading coefficient and hence takes negative values for x large enough.

Suppose A is a common bound for all tuples $c = (c_1, \dots, c_k)$ satisfying $c_1 + \cdots + c_k = b + 1$ (note that there are only finitely many such tuples). Then, for any tuple $c' = (c'_1, \dots, c'_k)$ with $c'_1 + \cdots + c'_k > b$, there exists a tuple $c = (c_1, \dots, c_k)$ with $c_1 + \cdots + c_k = b + 1$ and $c'_i \geq c_i$, and then the inequality for c' follows from the inequality for c .

We can show part 2 either in a similar way, or by using that there are only finitely many such tuples. \square

Take A satisfying the assertion of Claim 2, and take N such that $n \geq N$ implies $a_n \geq A$. Then for each $n \geq N$, we have

$$(a_{n+1} - a_n) + \cdots + (a_{n+k} - a_n) = b.$$

By taking the difference of this equality and the equality for $n + 1$, we obtain

$$a_{n+k+1} - a_{n+1} = k(a_{n+1} - a_n)$$

for every $n \geq N$.

We conclude using an extremal principle. Let $d = \min\{a_{n+1} - a_n \mid n \geq N\}$, and suppose it is attained at some index $n \geq N$. Since

$$kd = k(a_{n+1} - a_n) = a_{n+k+1} - a_{n+1} = \sum_{i=1}^k (a_{n+i+1} - a_{n+i})$$

and each summand is at least d , we conclude that d is also attained at $n + 1, \dots, n + k$, and inductively at all $n' \geq n$. We see that the equation $P(x) = (x + d)(x + 2d) \cdots (x + kd)$ is true for infinitely many values of x (all $a_{n'}$ for $n' \geq n$), hence this is an equality of polynomials. Finally we use (backward) induction to show that $a_{n+1} - a_n = d$ for every $n \geq 1$.

Solution 2. We assume that (a_n) is not eventually constant. In this solution, we first prove an alternative version of Claim 1.

Claim 3. There exist infinitely many $n \geq 1$ with

$$a_n \leq \min\{a_{n+1}, \dots, a_{n+k}\}.$$

Proof. Suppose not, then for all but finitely many $n \geq 1$, it holds that $a_n > \min\{a_{n+1}, \dots, a_{n+k}\}$. Hence for all large enough n , there always exist some $1 \leq l \leq k$ such that $a_n > a_{n+l}$. This induces an infinite decreasing sequence $a_n > a_{n+l_1} > a_{n+l_2} > \dots$ of positive integers, which is absurd. \square

We use Claim 3 to quickly settle the case $P(x) = x^k$. In that case, for every n with $a_n \leq \min\{a_{n+1}, \dots, a_{n+k}\}$, since $a_{n+1} \cdots a_{n+k} = a_n^k$, it implies $a_n = a_{n+1} = \dots = a_{n+k}$. This shows that the sequence is eventually constant, which contradicts our assumption.

From now on, assume

$$P(x) > x^k \text{ for all } x > 0.$$

Claim 4. For every $M > 0$, there exists some $N > 0$ such that $a_n > M$ for all $n > N$.

Proof. Suppose there exists some $M > 0$, such that $a_n \leq M$ for infinitely many n . For each i with $a_i \leq M$, we consider the k -tuple $(a_{i+1}, \dots, a_{i+k})$. Then each of the terms in the k -tuple is bounded from above by $P(a_i)$, and hence by $P(M)$ too. Since the number of such k -tuples is bounded by $P(M)^k$, we deduce by the Pigeonhole Principle that there exist some indices $i < j$ such that $(a_{i+1}, \dots, a_{i+k}) = (a_{j+1}, \dots, a_{j+k})$. Since a_n is uniquely determined by the k terms before it, we conclude that $a_{i+k+1} = a_{j+k+1}$ must hold, and similarly $a_{i+l} = a_{j+l}$ for all $l \geq 0$, so the sequence is eventually periodic, for some period $p = j - i$.

Take K such that $a_n = a_{n+p}$ for every $n \geq K$. Then, by taking the products of the inequalities

$$a_n^k < P(a_n) = a_{n+1} \cdots a_{n+k}$$

for $K \leq n \leq K + p - 1$, we obtain

$$\begin{aligned} \prod_{n=K}^{K+p-1} a_n^k &< \prod_{n=K}^{K+p-1} a_{n+1} \cdots a_{n+k} \\ &= a_{K+1} a_{K+2}^2 \cdots a_{K+k-1}^{k-1} \left(\prod_{n=K+k}^{K+p} a_n \right)^k a_{K+p+1}^{k-1} \cdots a_{K+p+k-2}^2 a_{K+p+k-1} \\ &= \left(\prod_{n=K}^{K+p-1} a_n \right)^k \quad (\text{by periodicity}), \end{aligned}$$

which is a contradiction. \square

Write $P(x) = x^k + bx^{k-1} + Q(x)$, where $Q(x)$ is of degree at most $k-2$. Take M such that $x > M$ implies $x^{k-1} > Q(x)$.

Claim 5. There exist non-negative integers b_1, \dots, b_k such that $P(x) = (x + b_1) \cdots (x + b_k)$, and such that, for infinitely many $n \geq 1$, we have $a_{n+i} = a_n + b_i$ for every $1 \leq i \leq k$.

Proof. By Claims 3 and 4, there are infinitely many n such that

$$a_n > M \text{ and } a_n \leq \min\{a_{n+1}, \dots, a_{n+k}\}.$$

Call such indices n to be *good*. We claim that if n is a good index then

$$\max\{a_{n+1}, \dots, a_{n+k}\} \leq a_n + b.$$

Indeed, if $a_{n+i} \geq a_n + b + 1$, then together with $a_n \leq \min\{a_{n+1}, \dots, a_{n+k}\}$ and $a_n^{k-1} > Q(a_n)$, we have

$$a_n^k + (b+1)a_n^{k-1} > a_n^k + ba_n^{k-1} + Q(a_n) = P(a_n) \geq (a_n + b + 1)a_n^{k-1},$$

a contradiction.

Hence for each good index n , we may write $a_{n+i} = a_n + b_i$ for all $1 \leq i \leq k$ for some choices of (b_1, \dots, b_k) (which may depend on n) and $0 \leq b_i \leq b$. Again by Pigeonhole Principle, some k -tuple (b_1, \dots, b_k) must be chosen for infinitely such good indices n . This means that the equation $P(a_n) = (a_n + b_1) \cdots (a_n + b_k)$ is satisfied by infinitely many good indices n . By Claim 4, a_n is unbounded among these a_n 's, hence $P(x) = (x + b_1) \cdots (x + b_k)$ must hold identically. \square

Claim 6. We have $b_i = ib_1$ for all $1 \leq i \leq k$.

Proof. Call an index n *excellent* if $a_{n+i} = a_n + b_i$ for every $1 \leq i \leq k$. From Claim 5 we know there are infinitely many excellent n .

We first show that for any pair $1 \leq i < j \leq k$ there is $1 \leq l \leq k$ such that $b_j = b_i + b_l$. Indeed, for such i and j and for excellent n , $a_n + b_j$ (which is equal to a_{n+j}) divides $P(a_{n+i}) = \prod_{l=1}^k (a_n + b_i + b_l)$, and hence divides $\prod_{l=1}^k (b_i + b_l - b_j)$. Since $a_n + b_j$ is unbounded among excellent n , we have $\prod_{l=1}^k (b_i + b_l - b_j) = 0$, hence there is l such that $b_j = b_i + b_l$.

In particular, $b_j = b_i + b_l \geq b_i$, i.e. (b_1, \dots, b_k) is non-decreasing.

Suppose $b_1 = 0$ and n is an excellent number. In particular, it holds that $a_n = a_{n+1}$. Moreover, since

$$a_{n+k+1}P(a_n) = a_{n+1} \cdots a_{n+k+1} = a_{n+1}P(a_{n+1}),$$

we have $a_n = a_{n+1} = a_{n+k+1}$, which divides $P(a_{n+i}) = \prod_{l=1}^k (a_n + b_i + b_l)$ for each $1 \leq i \leq k$. Hence a_n divides $\prod_{l=1}^k (b_i + b_l)$. By the same reasoning, we have $b_i + b_l = 0$ for some l , but since $b_i, b_l \geq 0$ we obtain $b_i = 0$ for each $1 \leq i \leq k$.

Now suppose $b_1 \geq 1$. Then, for each $1 \leq i < j \leq k$, we have $b_j - b_i = b_l \geq b_1 \geq 1$, hence (b_1, \dots, b_k) is strictly increasing. Therefore, the $k-1$ elements $b_2 < b_3 < \dots < b_k$ are exactly equal to $b_1 + b_1 < \dots < b_1 + b_{k-1}$, since they cannot be equal to $b_1 + b_k$. This gives $b_i = ib_1$ for all $1 \leq i \leq k$ as desired. \square

Claim 6 implies $P(x) = (x + d)(x + 2d) \cdots (x + kd)$ for some $d \geq 1$, and there are infinitely many indices n with $a_{n+i} = a_n + id$ for $1 \leq i \leq k$. By backwards induction, $P(a_{n-1}) = a_n \cdots a_{n+k-1}$ implies $a_{n-1} = a_n - d$, and so on. Thus a_1, \dots, a_n forms an arithmetic progression with common difference d . Since n can be arbitrarily large, the whole sequence is an arithmetic progression too, as desired.

Comment 1. A typical solution would first show some kind of increasing property (assuming a_n is not constant), and then use that property to deduce informations on the numbers $a_{n+i} - a_n$ ($1 \leq i \leq k$) and/or on the polynomial P .

Solution 1 shows a strict one: $a_n < a_{n+1}$ (arguments after Claim 1), which makes the latter part easier. Solution 2 (Claims 3 and 4) shows only weaker increasing properties, which require more complicated/tricky arguments in the latter part but still can solve the problem.

Comment 2. It would be interesting to sort out the case when P can take negative integer coefficients, or (a_n) is just an integer sequence. Then a decreasing arithmetic progression is possible too, yet that is not the only possibility. There exist bounded examples such as $1, -1, 1, -1, \dots$ with $P(x) = -x^2$, or $0, 1, -1, 0, 1, -1, \dots$ with $P(x) = x^2 - 1$. If furthermore P is allowed to be non-monic, then the situation is even more unclear. For instance, the sequence $1, 2, 4, 8, \dots$ works for the polynomial $P(x) = 8x^2$.

A7. Let N be a positive integer. Prove that there exist three permutations a_1, a_2, \dots, a_N ; b_1, b_2, \dots, b_N ; and c_1, c_2, \dots, c_N of $1, 2, \dots, N$ such that

$$\left| \sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} - 2\sqrt{N} \right| < 2023$$

for every $k = 1, 2, \dots, N$.

(China)

Solution 1. The idea is to approximate the numbers $\sqrt{1}, \sqrt{2}, \dots, \sqrt{N}$ by the nearest integer with errors < 0.5 . This gives the following sequence

$$1, 1, 2, 2, 2, 2, 3, 3, 3, 3, 3, 3, 4, \dots$$

More precisely, for each $k \geq 1$, we round $\sqrt{k^2 - k + 1}, \dots, \sqrt{k^2 + k}$ to k , so that there are $2k$ copies of k .

Step 1. We first consider the easier case when N has the form

$$N = m(m+1).$$

In this case, the numbers $\sqrt{1}, \sqrt{2}, \dots, \sqrt{N}$ are approximated by the elements of the multiset $\{1_{\times 2}, 2_{\times 4}, 3_{\times 6}, \dots, m_{\times 2m}\}$. Let T_m denote “half of” the multiset, i.e.

$$T_m := \{1_{\times 1}, 2_{\times 2}, 3_{\times 3}, \dots, m_{\times m}\}.$$

We will prove by induction that there exists three permutations (u_k) , (v_k) , and (w_k) of the elements in the multiset T_m such that $u_k + v_k + w_k = 2m + 1$ is constant for $k = 1, 2, \dots, \frac{m(m+1)}{2}$.

When $m = 1$, take $1 + 1 + 1 = 3$. When $m = 2$, take $(1, 2, 2) + (2, 1, 2) + (2, 2, 1) = (5, 5, 5)$. Suppose that we have constructed three permutations (u_k) , (v_k) , and (w_k) of T_{m-1} satisfying $u_k + v_k + w_k = 2m - 1$ for every $k = 1, 2, \dots, \frac{m(m-1)}{2}$. For T_m , we note that

$$T_m = T_{m-1} \sqcup \{m_{\times m}\},$$

and also

$$T_m = (T_{m-1} + 1) \sqcup \{1, 2, \dots, m\}. \quad (1)$$

Here $T_{m-1} + 1$ means to add 1 to all elements in T_{m-1} . We construct the permutations (u'_k) , (v'_k) , and (w'_k) of T_m as follows:

- For $k = 1, 2, \dots, \frac{m(m-1)}{2}$, we set $u'_k = u_k$, $v'_k = v_k + 1$, $w'_k = w_k + 1$.
- For $k = \frac{m(m-1)}{2} + r$ with $r = 1, 2, \dots, m$, we set $u'_k = m$, $v'_k = r$, $w'_k = m + 1 - r$.

It is clear from (1) that (u'_k) , (v'_k) , and (w'_k) give three permutations of T_m , and that they satisfy $u'_k + v'_k + w'_k = 2m + 1$ for every $k = 1, 2, \dots, \frac{m(m+1)}{2}$.

The inductive construction can be visualised by the $3 \times \frac{m(m+1)}{2}$ matrix

$$\begin{bmatrix} u_1 & \dots & u_{m(m-1)/2} & m & \dots & m \\ v_1 + 1 & \dots & v_{m(m-1)/2} + 1 & 1 & \dots & m \\ w_1 + 1 & \dots & w_{m(m-1)/2} + 1 & m & \dots & 1 \end{bmatrix},$$

in which the three rows represent the permutations (u'_k) , (v'_k) , (w'_k) , and the sum of the three entries of each column is $2m + 1$.

Thus, when $N = m^2 + m$, we can construct permutations (a_k) , (b_k) , and (c_k) of $1, 2, \dots, N$ such that

$$2m + 1 - 1.5 < \sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} < 2m + 1 + 1.5. \quad (2)$$

This gives

$$\left| \sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} - 2\sqrt{N} \right| < 2.5 < 2023,$$

where we used that $-1 < 2m - 2\sqrt{m^2 + m} < 0$ for positive m .

Step 2. We now proceed to the general case. Let m be such that

$$m(m+1) \leq N < (m+1)(m+2).$$

Write $N = m(m+1) + t$ for some $t \in \{0, 1, \dots, 2m+1\}$ and let

$$L := \left\lfloor \frac{4}{9}N \right\rfloor.$$

We will make use of the following inequalities below:

$$N > m^2, \quad N < (m+2)^2, \quad t \leq 2m+1, \quad L+1 > 4N/9, \quad L \leq 4N/9.$$

As above, we construct three permutations (a_k) , (b_k) , and (c_k) of $1, 2, \dots, m(m+1)$ satisfying (2). Now we construct the three required permutations (A_k) , (B_k) , and (C_k) of $1, 2, \dots, N$ as follows:

For $k = 1, 2, \dots, m(m+1)$, if $a_k \leq L$, take $A_k = a_k$, and if $a_k > L$, take $A_k = a_k + t$. For $k = m(m+1) + r$ with $r = 1, 2, \dots, t$, set $A_k = L + r$. Define the permutations (B_k) and (C_k) similarly. Now for $k = 1, 2, \dots, m(m+1)$, we show $0 \leq \sqrt{A_k} - \sqrt{a_k} \leq 2$. The lower bound is obvious. If $m \leq 1$, then $N \leq 5$ and hence $\sqrt{A_k} - \sqrt{a_k} \leq \sqrt{5} - \sqrt{1} \leq 2$. If $m \geq 2$, then

$$\sqrt{A_k} - \sqrt{a_k} = \frac{A_k - a_k}{\sqrt{A_k} + \sqrt{a_k}} \leq \frac{t}{2\sqrt{L+1}} \leq \frac{2m+1}{\frac{4}{3}m} \leq 2.$$

We have similar inequalities for (B_k) and (C_k) . Thus

$$2\sqrt{N} - 4.5 < 2m + 1 - 1.5 \leq \sqrt{A_k} + \sqrt{B_k} + \sqrt{C_k} \leq 2m + 1 + 1.5 + 6 < 2\sqrt{N} + 8.5.$$

For $k = m^2 + m + 1, \dots, m^2 + m + t$, we have

$$2\sqrt{N} < 3\sqrt{L+1} \leq \sqrt{A_k} + \sqrt{B_k} + \sqrt{C_k} \leq 3\sqrt{L+t} \leq \sqrt{4N+9t} < 2\sqrt{N} + 8.5.$$

To sum up, we have defined three permutations (A_k) , (B_k) , and (C_k) of $1, 2, \dots, N$, such that

$$\left| \sqrt{A_k} + \sqrt{B_k} + \sqrt{C_k} - 2\sqrt{N} \right| < 8.5 < 2023.$$

holds for every $k = 1, 2, \dots, N$. □

Solution 2. This is a variation of Solution 1 that uses induction for Step 2.

Let n be an integer satisfying $0 \leq n \leq m+1$ and define the multiset $T_{m,n}$ by

$$T_{m,n} := \{1_{\times 1}, 2_{\times 2}, 3_{\times 3}, \dots, m_{\times m}, (m+1)_{\times n}\}.$$

In other words, $T_{m,0} = T_m$, $T_{m,n} = T_m \sqcup \{(m+1)_{\times n}\}$ and $T_{m,m+1} = T_{m+1}$, where T_m is the set defined in Solution 1.

Claim. There exist three permutations $(u_k), (v_k), (w_k)$ of $T_{m,n}$ such that

$$\begin{cases} u_k + v_k + w_k = 2m + 1 & (n = 0), \\ u_k + v_k + w_k \in \{2m + 1, 2m + 2, 2m + 3\} & (1 \leq n \leq m), \\ u_k + v_k + w_k = 2m + 3 & (n = m + 1). \end{cases}$$

Proof. We proceed by induction on m . If $n = 0$ or $n = m + 1$, the assertion can be proved as in Solution 1. If $1 \leq n \leq m$, we note that

$$T_{m,n} = T_{m-1,n} \sqcup \{m_{\times(m-n)}, (m+1)_{\times n}\} = (T_{m-1,n} + 1) \sqcup \{1, 2, \dots, m\}.$$

From the hypothesis of induction, it follows that we have three permutations $(u_k), (v_k), (w_k)$ of $T_{m-1,n}$ satisfying $u_k + v_k + w_k \in \{2m - 1, 2m, 2m + 1\}$ for every k . We construct the permutations $(u'_k), (v'_k)$, and (w'_k) of $T_{m,n}$ as follows:

- For $k = 1, 2, \dots, \frac{m(m-1)}{2} + n$, we set $u'_k = u_k$, $v'_k = v_k + 1$, and $w'_k = w_k + 1$.
- For $k = \frac{m(m-1)}{2} + n + r$ with $r = 1, 2, \dots, m$, we set $u'_k = m$ if $1 \leq r \leq m - n$ while $u'_k = m + 1$ if $m - n + 1 \leq r \leq m$, $v'_k = r$, and $w'_k = m + 1 - r$.

It is clear from the construction that $(u'_k), (v'_k)$, and (w'_k) give three permutations of $T_{m,n}$, and they satisfy $u'_k + v'_k + w'_k \in \{2m + 1, 2m + 2, 2m + 3\}$ for every $k = 1, 2, \dots, \frac{m(m+1)}{2} + n$.

Again, we can visualise the construction using the matrix

$$\begin{bmatrix} u_1 & \dots & u_{m(m-1)/2+n} & m & \dots & m & m+1 & \dots & m+1 \\ v_1 + 1 & \dots & v_{m(m-1)/2+n} + 1 & 1 & \dots & \dots & \dots & \dots & m \\ w_1 + 1 & \dots & w_{m(m-1)/2+n} + 1 & m & \dots & \dots & \dots & \dots & 1 \end{bmatrix}. \quad \square$$

In general, we have $m(m+1) \leq N < (m+1)(m+2)$ for some $m \geq 0$. Set $N = m(m+1) + t$ for some $t \in \{0, 1, \dots, 2m+1\}$. Then the approximation of $\{\sqrt{1}, \sqrt{2}, \dots, \sqrt{N}\}$ by the nearest integer with errors < 0.5 is a multiset

$$\{1_{\times 2}, 2_{\times 4}, \dots, m_{\times 2m}, (m+1)_{\times t}\} = T_{m,n_1} \sqcup T_{m,n_2}$$

with $n_1 = \lfloor t/2 \rfloor$ and $n_2 = \lceil t/2 \rceil$.

Since $0 \leq n_1 \leq n_2 \leq m+1$, by using the Claim we can construct permutations $(a_k), (b_k)$, and (c_k) to satisfy the following inequality:


$$2m + 1 - 1.5 < \sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} < 2m + 3 + 1.5.$$

Since $m < \sqrt{N} < m + 2$, it follows that

$$2\sqrt{N} - 4.5 < 2m + 1 - 1.5 < \sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} < 2m + 3 + 1.5 < 2\sqrt{N} + 4.5,$$

and so

$$\left| \sqrt{A_k} + \sqrt{B_k} + \sqrt{C_k} - 2\sqrt{N} \right| < 4.5 < 2023. \quad \square$$



Solution 3. This solution is based on the geometrical insight of equilateral triangles.

Step 1. We first consider the easier case of triangle numbers

$$N = \frac{m(m+1)}{2}.$$

As shown in the following picture, consider the triangular shaped lattice points inside an equilateral triangle ABC with a total of N points. The lattice is built in a way that the ℓ^{th} row has exactly ℓ points for each $\ell = 1, 2, \dots, m$. Rows are numbered in three different ways, one for each vertex.

Each point P_k in the triangular lattice is labelled with a triple of integers (a_k, b_k, c_k) as follows. The first coordinate is called the A -coordinate, and so on for B, C . To define the A -coordinate, denoted $W_a(\bullet)$, first label the lattice points by $1, 2, 3, \dots$ starting with the point closest to A and then going down the rows with the rule that within a row, the labelling is from left to right (see right picture). The B -coordinate, denoted $W_b(\bullet)$, is defined by rotating the A -coordinate counterclockwise by 120° . The C -coordinate, denoted $W_c(\bullet)$, similarly, by rotating the A -coordinate counterclockwise by 240° .

Assume that a point P lies in the ℓ_a^{th} row from the vertex A , in the ℓ_b^{th} row from the vertex B , and in the ℓ_c^{th} row from the vertex C . Note that ℓ_a is proportional to the height of A in the triangle, minus the height of P . Since inside an equilateral triangle, the sum of the lengths of the heights from a point to the three sides is independent of the point, we must have

$$\ell_a + \ell_b + \ell_c = 2m + 1 = \sqrt{8N + 1}.$$

Since there are exactly $1 + 2 + \dots + \ell = \frac{\ell(\ell+1)}{2}$ points in the first ℓ rows, the A -labeling $W_a(P)$ of the point P satisfies

$$\frac{\ell_a(\ell_a - 1)}{2} + 1 \leq W_a(P) \leq \frac{\ell_a(\ell_a + 1)}{2}.$$

In particular,

$$\left(\ell_a - \frac{1}{2}\right)^2 < 2W_a(P) < \left(\ell_a + \frac{1}{2}\right)^2.$$

Taking the cyclic sum gives

$$\left| \sqrt{2W_a(P)} + \sqrt{2W_b(P)} + \sqrt{2W_c(P)} - (\ell_a + \ell_b + \ell_c) \right| < \frac{3}{2}$$

and thus

$$\left| \sqrt{W_a(P)} + \sqrt{W_b(P)} + \sqrt{W_c(P)} - 2\sqrt{N + \frac{1}{8}} \right| < \frac{3}{2} \cdot \frac{1}{\sqrt{2}} = \frac{3\sqrt{2}}{4}.$$

Step 2. Now, for a general positive integer N , there exists a positive integer m such that

$$\frac{m(m-1)}{2} + 1 \leq N \leq \frac{m(m+1)}{2}.$$

Write $N = \frac{m(m+1)}{2} - t$ with $t \in \{0, 1, \dots, m-1\}$. We modify the above construction for $\frac{m(m+1)}{2}$ points into a construction for N points as follows. We remove t arbitrary points from the m^{th} row (namely the bottom row) of the triangular lattice. The remaining triangular lattice has $\frac{m(m+1)}{2} - t = N$ points, and we assign their A -, B -, and C -coordinates as before (in the same order, yet skipping over the points that are removed so that the coordinates exactly form permutations of $1, 2, \dots, N$).

For each point P in the triangular lattice (that was not removed earlier), suppose that it is in the ℓ_a^{th} , ℓ_b^{th} , and ℓ_c^{th} row when viewed from A , B , and C , respectively. Now the A -coordinates $W_a(P)$ still satisfies

$$\frac{\ell_a(\ell_a - 1)}{2} + 1 \leq W_a(P) \leq \frac{\ell_a(\ell_a + 1)}{2}.$$

The B -coordinate $W_b(P)$ satisfies

$$\frac{(\ell_b - 1)(\ell_b - 2)}{2} + 1 \leq W_b(P) \leq \frac{\ell_b(\ell_b + 1)}{2},$$

because, viewing from point B , we have removed either 0 or 1 point from each row, and the first $\ell_b - 1$ rows have at least $0 + 1 + \dots + (\ell_b - 2) = \frac{(\ell_b - 1)(\ell_b - 2)}{2}$ points left. For the same reason, the C -labeling $W_c(P)$ satisfies

$$\frac{(\ell_c - 1)(\ell_c - 2)}{2} + 1 \leq W_c(P) \leq \frac{\ell_c(\ell_c + 1)}{2}.$$

From this, we deduce that

$$\begin{aligned} \ell_a - \frac{1}{2} &< \sqrt{2W_a(P)} < \ell_a + \frac{1}{2}, \\ \ell_b - \frac{3}{2} &< \sqrt{2W_b(P)} < \ell_b + \frac{1}{2}, \\ \ell_c - \frac{3}{2} &< \sqrt{2W_c(P)} < \ell_c + \frac{1}{2}. \end{aligned}$$

Combining all above with the inequalities $2m-1 < 2\sqrt{2N} < 2m+1$ and $\ell_a + \ell_b + \ell_c = 2m+1$, we deduce that

$$\begin{aligned} 2\sqrt{2N} - \frac{7}{2} &< (2m+1) - \frac{7}{2} < \sqrt{2W_a(P)} + \sqrt{2W_b(P)} + \sqrt{2W_c(P)} \\ &< (2m+1) + \frac{3}{2} < 2\sqrt{2N} + \frac{7}{2}. \end{aligned}$$

Therefore, for each point P , we have

$$|\sqrt{W_a(P)} + \sqrt{W_b(P)} + \sqrt{W_c(P)} - 2\sqrt{N}| < \frac{7}{2} \cdot \frac{1}{\sqrt{2}} < 2.5 < 2013.$$

We may finally order of the N points in an arbitrary way. Then the A -labelings $W_a(\bullet)$ give the permutation a_1, \dots, a_N , the B -labelings $W_b(\bullet)$ give b_1, \dots, b_N , and the C -labelings $W_c(\bullet)$ give c_1, \dots, c_N .

For each $k = 1, 2, \dots, N$, we have

$$|\sqrt{a_k} + \sqrt{b_k} + \sqrt{c_k} - 2\sqrt{N}| < 2.5 < 2023.$$

□

Comment. We can make the same argument as in Solution 3 without using geometry or diagram instead using barycentric coordinates in integers and lexicographic order.

For $N = \frac{m(m+1)}{2}$, consider a set

$$X = \{(x, y, z) \in \mathbb{Z}^3 \mid 0 \leq x, y, z \leq m-1, x+y+z = m-1\}.$$

and the lexicographic order of X , i.e.

$$(x, y, z) > (x', y', z') \iff \begin{cases} x > x' \text{ or} \\ x = x' \text{ and } y > y' \text{ or} \\ x = x' \text{ and } y = y' \text{ and } z > z'. \end{cases}$$

Then for an element $Q_k = (x_k, y_k, z_k)$

- Define $W_a(Q_k)$ so that (x_k, y_k, z_k) is the $W_a(Q_k)^{\text{th}}$ biggest element in X .
- Define $W_b(Q_k)$ so that (y_k, z_k, x_k) is the $W_b(Q_k)^{\text{th}}$ biggest element in $X' = \{(y, z, x) \mid (x, y, z) \in X\}$.
- Define $W_c(Q_k)$ so that (z_k, x_k, y_k) is the $W_c(Q_k)^{\text{th}}$ biggest element in $X'' = \{(z, x, y) \mid (x, y, z) \in X\}$.

The same argument as in Solution 2 then holds.

Observe that for an element $Q_k = (x_k, y_k, z_k)$, it holds that $\ell_a = m - x_k$, $\ell_b = m - y_k$, and $\ell_c = m - z_k$.

Combinatorics

C1. Let m and n be positive integers greater than 1. In each unit square of an $m \times n$ grid lies a coin with its tail-side up. A *move* consists of the following steps:

1. select a 2×2 square in the grid;
2. flip the coins in the top-left and bottom-right unit squares;
3. flip the coin in either the top-right or bottom-left unit square.

Determine all pairs (m, n) for which it is possible that every coin shows head-side up after a finite number of moves.

(Thailand)

Answer: The answer is all pairs (m, n) satisfying $3 \mid mn$.

Solution 1. Let us denote by (i, j) -square the unit square in the i^{th} row and the j^{th} column.


We first prove that when $3 \mid mn$, it is possible to make all the coins show head-side up. For integers $1 \leq i \leq m-1$ and $1 \leq j \leq n-1$, denote by $A(i, j)$ the move that flips the coin in the (i, j) -square, the $(i+1, j+1)$ -square and the $(i, j+1)$ -square. Similarly, denote by $B(i, j)$ the move that flips the coin in the (i, j) -square, $(i+1, j+1)$ -square, and the $(i+1, j)$ -square. Without loss of generality, we may assume that $3 \mid m$.

Case 1: n is even.

We apply the moves

- $A(3k-2, 2l-1)$ for all $1 \leq k \leq \frac{m}{3}$ and $1 \leq l \leq \frac{n}{2}$,
- $B(3k-1, 2l-1)$ for all $1 \leq k \leq \frac{m}{3}$ and $1 \leq l \leq \frac{n}{2}$.

This process will flip each coin exactly once, hence all the coins will face head-side up afterwards.



Case 2: n is odd.

We start by applying

- $A(3k-2, 2l-1)$ for all $1 \leq k \leq \frac{m}{3}$ and $1 \leq l \leq \frac{n-1}{2}$,
- $B(3k-1, 2l-1)$ for all $1 \leq k \leq \frac{m}{3}$ and $1 \leq l \leq \frac{n-1}{2}$

as in the previous case. At this point, the coins on the rightmost column have tail-side up and the rest of the coins have head-side up. We now apply the moves

- $A(3k-2, n-1)$, $A(3k-1, n-1)$ and $B(3k-2, n-1)$ for every $1 \leq k \leq \frac{m}{3}$.

For each k , the three moves flip precisely the coins in the $(3k - 2, n)$ -square, the $(3k - 1, n)$ -square, and the $(3k, n)$ -square. Hence after this process, every coin will face head-side up.

We next prove that mn being divisible by 3 is a necessary condition. We first label the (i, j) -square by the remainder of $i + j - 2$ when divided by 3, as shown in the figure.

0	1	2	0	...
1	2	0	1	...
2	0	1	2	...
0	1	2	0	...
\vdots	\vdots	\vdots	\vdots	\ddots

Let $T(c)$ be the number of coins facing head-side up in those squares whose label is c . The main observation is that each move does not change the parity of both $T(0) - T(1)$ and $T(1) - T(2)$, since a move flips exactly one coin in a square with each label. Initially, all coins face tail-side up at the beginning, thus all of $T(0), T(1), T(2)$ are equal to 0. Hence it follows that any configuration that can be achieved from the initial state must satisfy the parity condition of

$$T(0) \equiv T(1) \equiv T(2) \pmod{2}.$$

We now calculate the values of T for the configuration in which all coins are facing head-side up.

- When $m \equiv n \equiv 1 \pmod{3}$, we have $T(0) - 1 = T(1) = T(2) = \frac{mn-1}{3}$.
- When $m \equiv 1 \pmod{3}$ and $n \equiv 2 \pmod{3}$, or $m \equiv 2 \pmod{3}$ and $n \equiv 1 \pmod{3}$, we have $T(0) - 1 = T(1) - 1 = T(2) = \frac{mn-2}{3}$.
- When $m \equiv n \equiv 2 \pmod{3}$, we have $T(0) = T(1) - 1 = T(2) = \frac{mn-1}{3}$.
- When $m \equiv 0 \pmod{3}$ or $n \equiv 0 \pmod{3}$, we have $T(0) = T(1) = T(2) = \frac{mn}{3}$.

From this calculation, we see that $T(0)$, $T(1)$ and $T(2)$ has the same parity only when mn is divisible by 3.

Comment 1. The original proposal of the problem also included the following question as part (b):

For each pair (m, n) of integers greater than 1, how many configurations can be obtained by applying a finite number of moves?

An explicit construction of a sequence of moves shows that $T(0)$, $T(1)$, and $T(2)$ having the same parity is a necessary and sufficient condition for a configuration to obtainable after a finite sequence of moves, and this shows that the answer is 2^{mn-2} .

Comment 2. A significantly more difficult problem is to ask the following question: for pairs (m, n) such that the task is possible (i.e. $3 \mid mn$), what is the smallest number of moves required to complete this task? The answer is:

- $\frac{mn}{3}$ if mn is even;
- $\frac{mn}{3} + 2$ if mn is odd.

To show this, we observe that we can flip all coins in any 2×3 (or 3×2) by using a minimum of two moves. Furthermore, when mn is odd with $3 \mid mn$, it is impossible to tile an $m \times n$ table with one type of L-tromino and its 180° -rotated L-tromino (disallowing rotations and reflections). The only known proof of the latter claim is lengthy and difficult, and it requires some group-theoretic arguments by studying the tile homotopy group given by these two L-tromino tiles. This technique was developed by J. H. Conway and J. C. Lagarias in *Tiling with Polyominoes and Combinatorial Group Theory*, Journal of Combinatorial Group Theory, Series A **53**, 183-208 (1990).

Comment 3. Here is neat way of defining the invariant. Consider a finite field $\mathbb{F}_4 = \{0, 1, \omega, \omega + 1\}$, where $1 + 1 = \omega^2 + \omega + 1 = 0$ in \mathbb{F}_4 . Consider the set

$$H = \{(i, j) | 1 \leq i \leq m, 1 \leq j \leq n, \text{ the coin in the } (i, j)\text{-square is head-side up}\}.$$

and the invariant

$$I(H) = \sum_{(i,j) \in H} \omega^{i+j} \in \mathbb{F}_4.$$

Then the value of $I(H)$ does not change under applying moves, and when all coins are tail-side up, it holds that $I(H) = 0$. On the other hand, its value when all coins are head-side up can be computed as

$$I(H) = \sum_{i=1}^m \sum_{j=1}^n \omega^{i+j} = \left(\sum_{i=1}^m \omega^i \right) \left(\sum_{j=1}^n \omega^j \right).$$

This is equal to $0 \in \mathbb{F}_4$ if and only if $3 \mid mn$.

C2.

Determine the maximal length L of a sequence a_1, \dots, a_L of positive integers satisfying both the following properties:

- every term in the sequence is less than or equal to 2^{2023} , and
- there does not exist a consecutive subsequence a_i, a_{i+1}, \dots, a_j (where $1 \leq i \leq j \leq L$) with a choice of signs $s_i, s_{i+1}, \dots, s_j \in \{1, -1\}$ for which

$$s_i a_i + s_{i+1} a_{i+1} + \dots + s_j a_j = 0.$$

(Czech Republic)

Answer: The answer is $L = 2^{2024} - 1$.

Solution. We prove more generally that the answer is $2^{k+1} - 1$ when 2^{2023} is replaced by 2^k for an arbitrary positive integer k . Write $n = 2^k$.

We first show that there exists a sequence of length $L = 2n - 1$ satisfying the properties. For a positive integer x , denote by $v_2(x)$ the maximal nonnegative integer v such that 2^v divides x . Consider the sequence a_1, \dots, a_{2n-1} defined as

$$a_i = 2^{k-v_2(i)}.$$

For example, when $k = 2$ and $n = 4$, the sequence is

$$4, 2, 4, 1, 4, 2, 4.$$

This indeed consists of positive integers less than or equal to $n = 2^k$, because $0 \leq v_2(i) \leq k$ for $1 \leq i \leq 2^{k+1} - 1$.

Claim 1. This sequence a_1, \dots, a_{2n-1} does not have a consecutive subsequence with a choice of signs such that the signed sum equals zero.

Proof. Let $1 \leq i \leq j \leq 2n - 1$ be integers. The main observation is that amongst the integers

$$i, i+1, \dots, j-1, j,$$

there exists a unique integer x with the maximal value of $v_2(x)$. To see this, write $v = \max(v_2(i), \dots, v_2(j))$. If there exist at least two multiples of 2^v amongst $i, i+1, \dots, j$, then one of them must be a multiple of 2^{v+1} , which is a contradiction.

Therefore there is exactly one $i \leq x \leq j$ with $v_2(x) = v$, which implies that all terms except for $a_x = 2^{k-v}$ in the sequence

$$a_i, a_{i+1}, \dots, a_j$$

are a multiple of 2^{k-v+1} . The same holds for the terms $s_i a_i, s_{i+1} a_{i+1}, \dots, s_j a_j$, hence the sum cannot be equal to zero. \square

We now prove that there does not exist a sequence of length $L \geq 2n$ satisfying the conditions of the problem. Let a_1, \dots, a_L be an arbitrary sequence consisting of positive integers less than or equal to n . Define a sequence s_1, \dots, s_L of signs recursively as follows:

- when $s_1 a_1 + \dots + s_{i-1} a_{i-1} \leq 0$, set $s_i = +1$,
- when $s_1 a_1 + \dots + s_{i-1} a_{i-1} \geq 1$, set $s_i = -1$.

Write

$$b_i = \sum_{j=1}^i s_j a_j = s_1 a_1 + \dots + s_i a_i,$$

and consider the sequence

$$0 = b_0, b_1, b_2, \dots, b_L.$$

Claim 2. All terms b_i of the sequence satisfy $-n + 1 \leq b_i \leq n$.

Proof. We prove this by induction on i . It is clear that $b_0 = 0$ satisfies $-n + 1 \leq 0 \leq n$. We now assume $-n + 1 \leq b_{i-1} \leq n$ and show that $-n + 1 \leq b_i \leq n$.

Case 1: $-n + 1 \leq b_{i-1} \leq 0$.

Then $b_i = b_{i-1} + a_i$ from the definition of s_i , and hence

$$-n + 1 \leq b_{i-1} < b_{i-1} + a_i \leq b_{i-1} + n \leq n.$$

Case 2: $1 \leq b_{i-1} \leq n$.

Then $b_i = b_{i-1} - a_i$ from the definition of s_i , and hence

$$-n + 1 \leq b_{i-1} - n \leq b_{i-1} - a_i < b_{i-1} \leq n.$$

This finishes the proof. □


Because there are $2n$ integers in the closed interval $[-n + 1, n]$ and at least $2n + 1$ terms in the sequence b_0, b_1, \dots, b_L (as $L + 1 \geq 2n + 1$ by assumption), the pigeonhole principle implies that two distinct terms b_{i-1}, b_j (where $1 \leq i \leq j \leq L$) must be equal. Subtracting one from another, we obtain

$$s_i a_i + \dots + s_j a_j = b_j - b_{i-1} = 0$$

as desired.

Comment. The same argument gives a bound $L \leq 2n - 1$ that works for all n , but this bound is not necessarily sharp when n is not a power of 2. For instance, when $n = 3$, the longest sequence has length $L = 3$.

C3. Let n be a positive integer. We arrange $1 + 2 + \cdots + n$ circles in a triangle with n rows, such that the i^{th} row contains exactly i circles. The following figure shows the case $n = 6$.




In this triangle, a *ninja-path* is a sequence of circles obtained by repeatedly going from a circle to one of the two circles directly below it. In terms of n , find the largest value of k such that if one circle from every row is coloured red, we can always find a ninja-path in which at least k of the circles are red.

(Netherlands)

Answer: The maximum value is $k = 1 + \lfloor \log_2 n \rfloor$.


Solution 1. Write $N = \lfloor \log_2 n \rfloor$ so that we have $2^N \leq n \leq 2^{N+1} - 1$.

We first provide a construction where every ninja-path passes through at most $N + 1$ red circles. For the row $i = 2^a + b$ for $0 \leq a \leq N$ and $0 \leq b < 2^a$, we colour the $(2b + 1)^{\text{th}}$ circle.



Then every ninja-path passes through at most one red circle in each of the rows $2^a, 2^a + 1, \dots, 2^{a+1} - 1$ for each $0 \leq a \leq N$. It follows that every ninja-path passes through at most $N + 1$ red circles.

We now prove that for every colouring, there exists a ninja-path going through at least $N + 1$ red circles. For each circle C , we assign the maximum number of red circles in a ninja-path that starts at the top of the triangle and ends at C .



Note that

- if C is not red, then the number assigned to C is the maximum of the number assigned to the one or two circles above C , and
- if C is red, then the number assigned to C is one plus the above maximum.

Write v_1, \dots, v_i for the numbers in row i , and let v_m be the maximum among these numbers. Then the numbers in row $i + 1$ will be at least

$$v_1, \dots, v_{m-1}, v_m, v_m, v_{m+1}, \dots, v_i,$$

not taking into account the fact that one of the circles in row $i + 1$ is red. On the other hand, for the red circle in row $i + 1$, the lower bound on the assigned number can be increased by 1. Therefore the sum of the numbers in row $i + 1$ is at least

$$(v_1 + \dots + v_i) + v_m + 1.$$

Using this observation, we prove the following claim.

Claim 1. Let σ_k be the sum of the numbers assigned to circles in row k . Then for $0 \leq j \leq N$, we have $\sigma_{2^j} \geq j \cdot 2^j + 1$.

Proof. We use induction on j . This is clear for $j = 0$, since the number in the first row is always 1. For the induction step, suppose that $\sigma_{2^j} \geq j \cdot 2^j + 1$. Then the maximum value assigned to a circle in row 2^j is at least $j + 1$. As a consequence, for every $k \geq 2^j$, there is a circle on row k with number at least $j + 1$. Then by our observation above, we have

$$\sigma_{k+1} \geq \sigma_k + (j + 1) + 1 = \sigma_k + (j + 2).$$

Then we get

$$\sigma_{2^{j+1}} \geq \sigma_{2^j} + 2^j(j + 2) \geq j \cdot 2^j + 1 + 2^j(j + 2) = (j + j + 2)2^j + 1 = (j + 1)2^{j+1} + 1.$$

This completes the inductive step. \square


For $j = N$, this immediately implies that some circle in row 2^N has number at least $N + 1$. This shows that there is a ninja-path passing through at least $N + 1$ red circles.

Solution 2. We give an alternative proof that there exists a ninja-path passing through at least $N + 1$ red circles. Assign numbers to circles as in the previous solution, but we only focus on the numbers assigned to red circles.

For each positive integer i , denote by e_i the number of red circles with number i .

Claim 2. If the red circle on row l has number i , then $e_i \leq l$.

Proof. Note that if two circles C and C' are both assigned the same number i , then there cannot be a ninja-path joining the two circles. We partition the triangle into a smaller triangle with the red circle in row l at its top along with $l - 1$ lines that together cover all other circles.



In each set, there can be at most one red circle with number i , and therefore $e_i \leq l$. \square

We observe that if there exists a red circle C with number $i \geq 2$, then there also exists a red circle with number $i - 1$ in some row that is above the row containing C . This is because the second last red circle in the ninja-path ending at C has number $i - 1$.

Claim 3. We have $e_i \leq 2^{i-1}$ for every positive integer i .

Proof. We prove by induction on i . The base case $i = 1$ is clear, since the only red circle with number 1 is the one at the top of the triangle. We now assume that the statement is true for $1 \leq i \leq j-1$ and prove the statement for $i = j$. If $e_j = 0$, there is nothing to prove. Otherwise, let l be minimal such that the red circle on row l has number j . Then all the red circles on row $1, \dots, l-1$ must have number less than j . This shows that

$$l-1 \leq e_1 + e_2 + \dots + e_{j-1} \leq 1 + 2 + \dots + 2^{j-2} = 2^{j-1} - 1.$$


This proves that $l \leq 2^{j-1}$, and by Claim 2, we also have $e_j \leq l$. Therefore $e_j \leq 2^{j-1}$. \square

We now see that

$$e_1 + e_2 + \dots + e_N \leq 1 + \dots + 2^{N-1} = 2^N - 1 < n.$$

Therefore there exists a red circle with number at least $N+1$, which means that there exists a ninja-path passing through at least $N+1$ red circles.

Solution 3. We provide yet another proof that there exists a ninja-path passing through at least $N+1$ red circles. In this solution, we assign to a circle C the maximum number of red circles on a ninja-path *starting* at C (including C itself).



Denote by f_i the number of red circles with number i . Note that if a red circle C has number i , and there is a ninja-path from C to another red circle C' , then the number assigned to C' must be less than i .

Claim 4. If the red circle on row l has number less than or equal to i , then $f_i \leq l$.

Proof. This proof is same as the proof of Claim 2. The additional input is that if the red circle on row l has number strictly less than i , then the smaller triangle cannot have a red circle with number i . \square

Claim 5. We have

$$f_1 + f_2 + \dots + f_i \leq n - \left\lfloor \frac{n}{2^i} \right\rfloor$$

for all $0 \leq i \leq N$.

Proof. We use induction on i . The base case $i = 0$ is clear as the left hand side is the empty sum and the right hand side is zero. For the induction step, we assume that $i \geq 1$ and that the statement is true for $i-1$. Let l be minimal such that the red circle on row l has number less than or equal to i . Then all the red circles with number less than or equal to i lie on rows $l, l+1, \dots, n$, and therefore

$$f_1 + f_2 + \dots + f_i \leq n - l + 1.$$

On the other hand, the induction hypothesis together with the fact that $f_i \leq l$ shows that

$$f_1 + \dots + f_{i-1} + f_i \leq n - \left\lfloor \frac{n}{2^{i-1}} \right\rfloor + l.$$

Averaging the two inequalities gives

$$f_1 + \dots + f_i \leq n - \frac{1}{2} \left\lfloor \frac{n}{2^{i-1}} \right\rfloor + \frac{1}{2}.$$

Since the left hand side is an integer, we conclude that

$$f_1 + \cdots + f_i \leq n - \left\lfloor \frac{1}{2} \left\lfloor \frac{n}{2^{i-1}} \right\rfloor \right\rfloor = n - \left\lfloor \frac{n}{2^i} \right\rfloor.$$

This completes the induction step. □

Taking $i = N$, we obtain

$$f_1 + f_2 + \cdots + f_N \leq n - \left\lfloor \frac{n}{2^N} \right\rfloor < n.$$

This implies that there exists a ninja-path passing through at least $N + 1$ red circles.

Comment. Using essentially the same argument, one may inductively prove

$$e_a + e_{a+1} + \cdots + e_{a+i-1} \leq n - \left\lfloor \frac{n}{2^i} \right\rfloor.$$

instead. Taking $a = 1$ and $i = N$ gives the desired statement.

C4. Let $n \geq 2$ be a positive integer. Paul has a $1 \times n^2$ rectangular strip consisting of n^2 unit squares, where the i^{th} square is labelled with i for all $1 \leq i \leq n^2$. He wishes to cut the strip into several pieces, where each piece consists of a number of consecutive unit squares, and then *translate* (without rotating or flipping) the pieces to obtain an $n \times n$ square satisfying the following property: if the unit square in the i^{th} row and j^{th} column is labelled with a_{ij} , then $a_{ij} - (i + j - 1)$ is divisible by n .

Determine the smallest number of pieces Paul needs to make in order to accomplish this.

(U.S.A.)

Answer: The minimum number of pieces is $2n - 1$.

Solution 1. For the entirety of the solution, we shall view the labels as taking values in $\mathbb{Z}/n\mathbb{Z}$, as only their values modulo n play a role.

Here are two possible constructions consisting of $2n - 1$ pieces.

1. Cut into pieces of sizes $n, 1, n, 1, \dots, n, 1, 1$, and glue the pieces of size 1 to obtain the last row.
2. Cut into pieces of sizes $n, 1, n - 1, 2, n - 2, \dots, n - 1, 1$, and switch the pairs of consecutive strips that add up to size n .

We now prove that using $2n - 1$ pieces is optimal. It will be more helpful to think of the reverse process: start with n pieces of size $1 \times n$, where the k^{th} piece has squares labelled $k, k + 1, \dots, k + n - 1$. The goal is to restore the original $1 \times n^2$ strip.

Note that each piece, after cutting at appropriate places, is of the form $a, a + 1, \dots, b - 1$. Construct an (undirected but not necessarily simple) graph Γ with vertices labelled by $1, \dots, n$, where a piece of the form $a, a + 1, \dots, b - 1$ corresponds to an edge from a to b . We make the following observations.

- The cut pieces came from the k^{th} initial piece $k, k + 1, \dots, k + n - 1$ corresponds to a cycle γ_k (possibly of length 1) containing the vertex k .
- Since it is possible to rearrange the pieces into one single $1 \times n^2$ strip, the graph Γ has an Eulerian cycle.
- The number of edges of Γ is equal to the total number of cut pieces.

The goal is to prove that Γ has at least $2n - 1$ edges. Since Γ has an Eulerian cycle, it is connected. For every $1 \leq k \leq n$, pick one edge from γ_k , delete it from Γ to obtain a new graph Γ' . Since no two cycles γ_i and γ_j share a common edge, removing one edge from each cycle does not affect the connectivity of the graph. This shows that the new graph Γ' must also be connected. Therefore Γ' has at least $n - 1$ edges, which means that Γ has at least $2n - 1$ edges.

Solution 2. We provide an alternative proof that at least $2n - 1$ pieces are needed. Instead of having a linear strip, we work with a number of *circular strips*, each having length a multiple of n and labelled as

$$1, 2, \dots, n, 1, 2, \dots, n, \dots, 1, 2, \dots, n,$$

where there are n^2 cells in total across all circular strips. The goal is still to create the $n \times n$ square by cutting and translating. Here, when we say “translating” the strips, we imagine that each cell has a number written on it and the final $n \times n$ square is required to have every number written in the same upright, non-mirrored orientation.

Note that the number of cuts will be equal to the number of pieces, because performing $l \geq 1$ cuts on a single circular strip results in l pieces.

Consider any “seam” in the interior of the final square, between two squares S and T , so that S and T belongs to two separate pieces. We are interested in the positions of these two squares in the original circular strips, with the aim of removing the seam.

- If the two squares S and T come from the same circular strip and are adjacent, then the cut was unnecessary and we can simply remove the seam and reduce the number of required cuts by 1. The circular strips are not affected.
- If these two squares S and T were not adjacent, then they are next to two different cuts (either from the same circular strip or two different circular strips). Denote the two cuts by $(S|Y)$ and $(X|T)$. We perform these two cuts and then glue the pieces back according to $(S|T)$ and $(X|Y)$. Performing this move would either split one circular strip into two or merge two circular strips into one, changing the number of circular strips by at most one. Afterwards, we may eliminate cut $(S|T)$ since it is no longer needed, which also removes the corresponding seam from the final square.

By iterating this process, eventually we reach a state where there are some number of circular strips, but the final $n \times n$ square no longer has any interior seams.

Since no two rows of the square can be glued together while maintaining the consecutive numbering, the only possibility is to have exactly n circular strips, each with length n . In this state at least n cuts are required to reassemble the square. Recall that each seam removal operation changed the number of circular strips by at most one. So if we started with only one initial circular strip, then at least $n - 1$ seams were removed. Hence in total, at least $n + (n - 1) = 2n - 1$ cuts are required to transform one initial circular strip into the final square. Hence at least $2n - 1$ pieces are required to achieve the desired outcome.

Solution 3. As with the previous solution, we again work with *circular strips*. In particular, we start out with k *circular strips*, each having length a multiple of n and labelled as

$$1, 2, \dots, n, 1, 2, \dots, n, \dots, 1, 2, \dots, n,$$

where there are n^2 cells in total across all k circular strips. The goal is still to create the $n \times n$ square by cutting and translating the circular strips.

Claim. Constructing the $n \times n$ square requires at least $2n - k$ cuts (or alternatively, $2n - k$ pieces).

Proof. We prove by induction on n . The base case $n = 1$ is clear, because we can only have $k = 1$ and the only way of producing a 1×1 square from a 1×1 circular strip is by making a single cut. We now assume that $n \geq 2$ and the statement is true for $n - 1$.

Each cut is a cut between a cell of label i on the left and a cell of label $i + 1$ on the right side, for a unique $1 \leq i \leq n$. Let a_i be the number of such cuts, so that $a_1 + a_2 + \dots + a_n$ is the total number of cuts. Since all the left and right edges of the $n \times n$ square at the end must be cut, we have $a_i \geq 1$ for all $1 \leq i \leq n$.

If $a_i \geq 2$ for all i , then

$$a_1 + a_2 + \dots + a_n \geq 2n > 2n - k$$

and hence there is nothing to prove. We therefore assume that there exist some $1 \leq m \leq n$ for which $a_m = 1$. This unique cut must form the two ends of the linear strip

$$m + 1, m + 2, \dots, m - 1 + n, m + n$$

from the final product. There are two cases.

Case 1: The strip is a single connected piece.

In this case, the strip must have come from a single circular strip of length exactly n . We now remove this circular strip from of the cutting and pasting process. By definition of m , none of the edges between m and $m + 1$ are cut. Therefore we may pretend that all the adjacent pairs of cells labelled m and $m + 1$ are single cells. The induction hypothesis then implies that

$$a_1 + \cdots + a_{m-1} + a_{m+1} + \cdots + a_n \geq 2(n-1) - (k-1).$$

Adding back in a_m , we obtain

$$a_1 + \cdots + a_n \geq 2(n-1) - (k-1) + 1 = 2n - k.$$

Case 2: The strip is not a single connected piece.

Say the linear strip $m + 1, \dots, m + n$ is composed of $l \geq 2$ pieces C_1, \dots, C_l . We claim that if we cut the initial circular strips along both the left and right end points of the pieces C_1, \dots, C_l , and then remove them, the remaining part consists of at most $k + l - 2$ connected pieces (where some of them may be circular and some of them may be linear). This is because C_l, C_1 form a consecutive block of cells on the circular strip, and removing $l - 1$ consecutive blocks from k circular strips results in at most $k + (l - 1) - 1$ connected pieces.

Once we have the connected pieces that form the complement of C_1, \dots, C_l , we may glue them back at appropriate endpoints to form circular strips. Say we get k' circular strips after this procedure. As we are gluing back from at most $k + l - 2$ connected pieces, we see that

$$k' \leq k + l - 2.$$

We again observe that to get from the new circular strips to the $n - 1$ strips of size $1 \times n$, we never have to cut along the cell boundary between labels m and $m + 1$. Therefore the induction hypothesis applies, and we conclude that the total number of pieces is bounded below by

$$l + (2(n-1) - k') \geq l + 2(n-1) - (k + l - 2) = 2n - k.$$

This finishes the induction step, and therefore the statement holds for all n . \square

Taking $k = 1$ in the claim, we see that to obtain a $n \times n$ square from a circular $1 \times n^2$ strip, we need at least $2n - 1$ connected pieces. This shows that constructing the $n \times n$ square out of a linear $1 \times n^2$ strip also requires at least $2n - 1$ pieces.

C5.

Elisa has 2023 treasure chests, all of which are unlocked and empty at first. Each day, Elisa adds a new gem to one of the unlocked chests of her choice, and afterwards, a fairy acts according to the following rules:

- if more than one chests are unlocked, it locks one of them, or
- if there is only one unlocked chest, it unlocks all the chests.

Given that this process goes on forever, prove that there is a constant C with the following property: Elisa can ensure that the difference between the numbers of gems in any two chests never exceeds C , regardless of how the fairy chooses the chests to lock.

(Israel)

Solution 1. We will prove that such a constant C exists when there are n chests for n an odd positive integer. In fact we can take $C = n - 1$. Elisa's strategy is simple: place a gem in the chest with the fewest gems (in case there are more than one such chests, pick one arbitrarily).

For each integer $t \geq 0$, let $a_1^t \leq a_2^t \leq \dots \leq a_n^t$ be the numbers of gems in the n chests at the end of the t^{th} day. In particular, $a_1^0 = \dots = a_n^0 = 0$ and

$$a_1^t + a_2^t + \dots + a_n^t = t.$$

For each $t \geq 0$, there is a unique index $m = m(t)$ for which $a_m^{t+1} = a_m^t + 1$. We know that $a_j^t > a_{m(t)}^t$ for all $j > m(t)$, since $a_{m(t)}^t < a_{m(t)}^{t+1} \leq a_j^{t+1} = a_j^t$. Elisa's strategy also guarantees that if an index j is greater than the remainder of t when divided by n (i.e. the number of locked chests at the end of the t^{th} day), then $a_j^t \geq a_{m(t)}^t$, because some chest with at most a_j^t gems must still be unlocked at the end of the t^{th} day.

Recall that a sequence $x_1 \leq x_2 \leq \dots \leq x_n$ of real numbers is said to *majorise* another sequence $y_1 \leq y_2 \leq \dots \leq y_n$ of real numbers when for all $1 \leq k \leq n$ we have

$$x_1 + x_2 + \dots + x_k \leq y_1 + y_2 + \dots + y_k$$

and

$$x_1 + x_2 + \dots + x_n = y_1 + y_2 + \dots + y_n.$$

Our strategy for proving $a_n^t - a_1^t \leq n - 1$ is to inductively show that the sequence (a_i^t) is majorised by some other sequence (b_i^t) .

We define this other sequence (b_i^t) as follows. Let $b_k^0 = k - \frac{n+1}{2}$ for $1 \leq k \leq n$. As n is odd, this is a strictly increasing sequence of integers, and the sum of its terms is 0. Now define $b_i^t = b_i^0 + \lfloor \frac{t-i}{n} \rfloor + 1$ for $t \geq 1$ and $1 \leq i \leq n$. Thus for $t \geq 0$,

$$b_i^{t+1} = \begin{cases} b_i^t & \text{if } t+1 \not\equiv i \pmod{n}, \\ b_i^t + 1 & \text{if } t+1 \equiv i \pmod{n}. \end{cases}$$

From these properties it is easy to see that

- $b_1^t + b_2^t + \dots + b_n^t = t$ for all $t \geq 0$, and
- $b_i^t \leq b_{i+1}^t$ for all $t \geq 0$ and $1 \leq i \leq n-1$, with the inequality being strict if $t \not\equiv i \pmod{n}$.

Claim 1. For each $t \geq 0$, the sequence of integers $b_1^t, b_2^t, \dots, b_n^t$ majorises the sequence of integers $a_1^t, a_2^t, \dots, a_n^t$.

Proof. We use induction on t . The base case $t = 0$ is trivial. Assume $t \geq 0$ and that (b_i^t) majorises (a_i^t) . We want to prove the same holds for $t + 1$.

First note that the two sequences (b_i^{t+1}) and (a_i^{t+1}) both sum up to $t + 1$. Next, we wish to show that for $1 \leq k < n$, we have

$$b_1^{t+1} + b_2^{t+1} + \cdots + b_k^{t+1} \leq a_1^{t+1} + a_2^{t+1} + \cdots + a_k^{t+1}.$$

When $t + 1$ is replaced by t , the above inequality holds by the induction hypothesis. For the sake of contradiction, suppose k is the smallest index such that the inequality for $t + 1$ fails. Since the left hand side increases by at most 1 during the transition from t to $t + 1$, the inequality for $t + 1$ can fail only if all of the following occur:

- $b_1^t + b_2^t + \cdots + b_k^t = a_1^t + a_2^t + \cdots + a_k^t$,
- $t + 1 \equiv j \pmod{n}$ for some $1 \leq j \leq k$ (so that $b_j^{t+1} = b_j^t + 1$),
- $m(t) > k$ (so that $a_i^{t+1} = a_i^t$ for $1 \leq i \leq k$).

The first point and the minimality of k tell us that b_1^t, \dots, b_k^t majorises a_1^t, \dots, a_k^t as well (again using the induction hypothesis), and in particular $b_k^t \geq a_k^t$.

The second point tells us that the remainder of t when divided by n is at most $k - 1$, so $a_k^t \geq a_{m(t)}^t$ (by Elisa's strategy). But by the third point ($m(t) \geq k + 1$) and the non-decreasing property of a_i^t , we must have the equalities $a_k^t = a_{k+1}^t = a_{m(t)}^t$. On the other hand, $a_k^t \leq b_k^t < b_{k+1}^t$, with the second inequality being strict because $t \not\equiv k \pmod{n}$. We conclude that

$$b_1^t + b_2^t + \cdots + b_{k+1}^t > a_1^t + a_2^t + \cdots + a_{k+1}^t,$$

a contradiction to the induction hypothesis. □

This completes the proof as it implies

$$a_n^t - a_1^t \leq b_n^t - b_1^t \leq b_n^0 - b_1^0 = n - 1.$$

Comment 1. The statement is true even when n is even. In this case, we instead use the initial state

$$b_k^0 = \begin{cases} k - \frac{n}{2} - 1 & k \leq \frac{n}{2}, \\ k - \frac{n}{2} & k > \frac{n}{2}. \end{cases}$$

The same argument shows that $C = n$ works.

Comment 2. The constants $C = n - 1$ for odd n and $C = n$ for even n are in fact optimal. To see this, we will assume that the fairy always locks a chest with the minimal number of gems. Then at every point, if a chest is locked, any other chest with fewer gems will also be locked. Thus $m(t)$ is always greater than the remainder of t when divided by n . This implies that the quantities

$$I_k = a_1^t + \cdots + a_k^t - b_1^t - \cdots - b_k^t$$

for each $0 \leq k \leq n$, do not increase regardless of how Elisa acts. If Elisa succeeds in keeping $a_n^t - a_1^t$ bounded, then these quantities must also be bounded; thus they are eventually constant, say for $t \geq t_0$. This implies that for all $t \geq t_0$, the number $m(t)$ is equal to 1 plus the remainder of t when divided by n .

Claim 2. For $T \geq t_0$ divisible by n , we have

$$a_1^T < a_2^T < \cdots < a_n^T.$$

Proof. Suppose otherwise, and let j be an index for which $a_j^T = a_{j+1}^T$. We have $m(T + k - 1) = k$ for all $1 \leq k \leq n$. Then $a_j^{T+j} > a_{j+1}^{T+j}$, which gives a contradiction. \square

This implies $a_n^T - a_1^T \geq n - 1$, which already proves optimality of $C = n - 1$ for odd n . For even n , note that the sequence (a_i^T) has sum divisible by n , so it cannot consist of n consecutive integers. Thus $a_n^T - a_1^T \geq n$ for n even.

Solution 2. We solve the problem when 2023 is replaced with an arbitrary integer n . We assume that Elisa uses the following strategy:

At the beginning of the $(nt + 1)^{\text{th}}$ day, Elisa first labels her chests as C_1^t, \dots, C_n^t so that before she adds in the gem, the number of gems in C_i^t is less than or equal C_j^t for all $1 \leq i < j \leq n$. Then for days $nt + 1, nt + 2, \dots, nt + n$, she adds a gem to chest C_i^t , where i is chosen to be minimal such that C_i^t is unlocked.

Denote by c_i^t the number of gems in chest C_i^t at the beginning of the $(nt + 1)^{\text{th}}$ day, so that

$$c_1^t \leq c_2^t \leq \dots \leq c_n^t$$

by construction. Also, denote by δ_i^t the total number of gems added to chest C_i^t during days $nt + 1, \dots, nt + n$. We make the following observations.

- We have $c_1^0 = c_2^0 = \dots = c_n^0 = 0$.
- We have $c_1^t + \dots + c_n^t = nt$, since n gems are added every n days.
- The sequence (c_i^{t+1}) is a permutation of the sequence $(c_i^t + \delta_i^t)$ for all $t \geq 0$.
- We have $\delta_1^t + \dots + \delta_n^t = n$ for all $t \geq 0$.
- Since Elisa adds a gem to an unlocked chest C_i^t with i minimal, we have

$$\delta_1^t + \delta_2^t + \dots + \delta_k^t \geq k$$

for every $1 \leq k \leq n$ and $t \geq 0$.

We now define another sequence of sequences of integers as follows.

$$d_i^0 = 3n(i - \frac{n+1}{2}), \quad d_i^t = d_i^0 + t.$$

We observe that

$$d_1^t + \dots + d_n^t = c_1^t + \dots + c_n^t = nt.$$

Claim 3. For each $t \geq 0$, the sequence (d_i^t) majorises the sequence (c_i^t) .

Proof. We induct on t . For $t = 0$, this is clear as all the terms in the sequence (c_i^t) are equal. For the induction step, we assume that (d_i^t) majorises (c_i^t) . Given $1 \leq k \leq n - 1$, we wish to show that

$$d_1^{t+1} + \dots + d_k^{t+1} \leq c_1^{t+1} + \dots + c_k^{t+1}.$$

Case 1: $c_1^{t+1}, \dots, c_k^{t+1}$ is a permutation of $c_1^t + \delta_1^t, \dots, c_k^t + \delta_k^t$.

Since $d_1^t + \dots + d_k^t \leq c_1^t + \dots + c_k^t$ by the induction hypothesis, we have

$$\sum_{i=1}^k d_i^{t+1} = k + \sum_{i=1}^k d_i^t \leq k + \sum_{i=1}^k c_i^t \leq \sum_{i=1}^k (c_i^t + \delta_i^t) = \sum_{i=1}^k c_i^{t+1}.$$

Case 2: $c_1^{t+1}, \dots, c_k^{t+1}$ is not a permutation of $c_1^t + \delta_1^t, \dots, c_k^t + \delta_k^t$.

In this case, we have $c_i^t + \delta_i^t > c_j^t + \delta_j^t$ for some $i \leq k < j$. It follows that

$$c_k^t + n \geq c_i^t + n \geq c_i^t + \delta_i^t > c_j^t + \delta_j^t \geq c_j^t \geq c_{k+1}^t.$$

Using $d_k^t + 3n = d_{k+1}^t$ and the induction hypothesis, we obtain

$$\begin{aligned} \sum_{i=1}^k c_i^{t+1} &\geq \sum_{i=1}^k c_i^t > c_1^t + \dots + c_{k-1}^t + \frac{1}{2}c_k^t + \frac{1}{2}c_{k+1}^t - \frac{n}{2} = \frac{1}{2} \sum_{i=1}^{k-1} c_i^t + \frac{1}{2} \sum_{i=1}^{k+1} c_i^t - \frac{n}{2} \\ &\geq \frac{1}{2} \sum_{i=1}^{k-1} d_i^t + \frac{1}{2} \sum_{i=1}^{k+1} d_i^t - \frac{n}{2} = n + \sum_{i=1}^k d_i^t \geq k + \sum_{i=1}^k d_i^t = \sum_{i=1}^k d_i^{t+1}, \end{aligned}$$

This finishes the induction step. □

It follows that


$$c_n^t - c_1^t \leq d_n^t - d_1^t = 3n(n-1).$$

From day $nt+1$ to day $n(t+1)+1$, Elisa adds n gems, and therefore the difference may increase by at most n . This shows that the difference of the number of gems in two chests never exceeds $C = 3n(n-1) + n$.

C6.


Let N be a positive integer, and consider an $N \times N$ grid. A *right-down path* is a sequence of grid cells such that each cell is either one cell to the right of or one cell below the previous cell in the sequence. A *right-up path* is a sequence of grid cells such that each cell is either one cell to the right of or one cell above the previous cell in the sequence.

Prove that the cells of the $N \times N$ grid cannot be partitioned into less than N right-down or right-up paths. For example, the following partition of the 5×5 grid uses 5 paths.




(Canada)

Solution 1. We define a *good parallelogram* to be a parallelogram composed of two isosceles right-angled triangles glued together as shown below.



Given any partition into k right-down or right-up paths, we can find a corresponding packing of good parallelograms that leaves an area of k empty. Thus, it suffices to prove that we must leave an area of at least N empty when we pack good parallelograms into an $N \times N$ grid. This is actually equivalent to the original problem since we can uniquely recover the partition into right-down or right-up paths from the corresponding packing of good parallelograms.



We draw one of the diagonals in each cell so that it does not intersect any of the good parallelograms. Now, view these segments as mirrors, and consider a laser entering each of the $4N$ boundary edges (with starting direction being perpendicular to the edge), bouncing along these mirrors until it exits at some other edge. When a laser passes through a good parallelogram, its direction goes back to the original one after bouncing two times. Thus, if the final direction of a laser is perpendicular to its initial direction, it must pass through at least

one empty triangle. Similarly, if the final direction of a laser is opposite to its initial direction, it must pass through at least two empty triangles. Using this, we will estimate the number of empty triangles in the $N \times N$ grid.

We associate the starting edge of a laser with the edge it exits at. Then, the boundary edges are divided into $2N$ pairs. These pairs can be classified into three types:


- (1) a pair of a vertical and a horizontal boundary edge,
- (2) a pair of boundary edges from the same side, and
- (3) a pair of boundary edges from opposite sides.

Since the beams do not intersect, we cannot have one type (3) pair from vertical boundary edges and another type (3) pair from horizontal boundary edges. Without loss of generality, we may assume that we have t pairs of type (3) and they are all from vertical boundary edges. Then, out of the remaining boundary edges, there are $2N$ horizontal boundary edges and $2N - 2t$ vertical boundary edges. It follows that there must be at least t pairs of type (2) from horizontal boundary edges. We know that a laser corresponding to a pair of type (1) passes through at least one empty triangle, and a laser corresponding to a pair of type (2) passes through at least two empty triangles. Thus, as the beams do not intersect, we have at least $(2N - 2t) + 2 \cdot t = 2N$ empty triangles in the grid, leaving an area of at least N empty as required.

Solution 2. We apply an induction on N . The base case $N = 1$ is trivial. Suppose that the claim holds for $N - 1$ and prove it for $N \geq 2$.

Let us denote the path containing the upper left corner by P . If P is right-up, then every cell in P is in the top row or in the leftmost column. By the induction hypothesis, there are at least $N - 1$ paths passing through the lower right $(N - 1) \times (N - 1)$ subgrid. Since P is not amongst them, we have at least N paths.

Next, assume that P is right-down. If P contains the lower right corner, then we get an $(N - 1) \times (N - 1)$ grid by removing P and glueing the remaining two parts together. The main idea is to extend P so that it contains the lower right corner and the above procedure gives a valid partition of an $(N - 1) \times (N - 1)$ grid.



We inductively construct Q , which denotes an extension of P as a right-down path. Initially, $Q = P$. Let A be the last cell of Q , B be the cell below A , and C be the cell to the right of A (if they exist). Suppose that A is not the lower right corner, and that (*) both B and C do not belong to the same path as A . Then, we can extend Q as follows (in case we have two or more options, we can choose any one of them to extend Q).

1. If B belongs to a right-down path R , then we add the part of R , from B to its end, to Q .
2. If C belongs to a right-down path R , then we add the part of R , from C to its end, to Q .
3. If B belongs to a right-up path R which ends at B , then we add the part of R in the same column as B to Q .

4. If C belongs to a right-up path R which starts at C , then we add the part of R in the same row as C to Q .
5. Otherwise, B and C must belong to the same right-up path R . In this case, we add B and the cell to the right of B to Q .

Note that if B does not exist, then case (4) must hold. If C does not exist, then case (3) must hold.

It is easily seen that such an extension also satisfies the hypothesis (*), so we can repeat this construction to get an extension of P containing the lower right corner, denoted by Q . We show that this is a desired extension, i.e. the partition of an $(N-1) \times (N-1)$ grid obtained by removing Q and glueing the remaining two parts together consists of right-down or right-up paths.

Take a path R in the partition of the $N \times N$ grid intersecting Q . If the intersection of Q and R occurs in case (1) or case (2), then there exists a cell D in R such that the intersection of Q and R is the part of R from D to its end, so R remains a right-down path after removal of Q . Similarly, if the intersection of Q and R occurs in case (3) or case (4), then R remains a right-up path after removal of Q . If the intersection of Q and R occurs in case (5), then this intersection has exactly two adjacent cells. After the removal of these two cells (as we remove Q), R is divided into two parts that are glued into a right-up path.

Thus, we may apply the induction hypothesis to the resulting partition of an $(N-1) \times (N-1)$ grid, to find that it must contain at least $N-1$ paths. Since P is contained in Q and is not amongst these paths, the original partition must contain at least N paths.

C7.

The Imomi archipelago consists of $n \geq 2$ islands. Between each pair of distinct islands is a unique ferry line that runs in both directions, and each ferry line is operated by one of k companies. It is known that if any one of the k companies closes all its ferry lines, then it becomes impossible for a traveller, no matter where the traveller starts at, to visit all the islands exactly once (in particular, not returning to the island the traveller started at).

Determine the maximal possible value of k in terms of n .

(Ukraine)

Answer: The largest k is $k = \lfloor \log_2 n \rfloor$.

Solution. We reformulate the problem using graph theory. We have a complete graph K_n on n nodes (corresponding to islands), and we want to colour the edges (corresponding to ferry lines) with k colours (corresponding to companies), so that every Hamiltonian path contains all k different colours. For a fixed set of k colours, we say that an edge colouring of K_n is *good* if every Hamiltonian path contains an edge of each one of these k colours.

We first construct a good colouring of K_n using $k = \lfloor \log_2 n \rfloor$ colours.

Claim 1. Take $k = \lfloor \log_2 n \rfloor$. Consider the complete graph K_n in which the nodes are labelled by $1, 2, \dots, n$. Colour node i with colour $\min(\lfloor \log_2 i \rfloor + 1, k)$ (so the colours of the first nodes are $1, 2, 2, 3, 3, 3, 3, 4, \dots$ and the last $n - 2^{k-1} + 1$ nodes have colour k), and for $1 \leq i < j \leq n$, colour the edge ij with the colour of the node i . Then the resulting edge colouring of K_n is good.

Proof. We need to check that every Hamiltonian path contains edges of every single colour. We first observe that the number of nodes assigned colour k is $n - 2^{k-1} + 1$. Since $n \geq 2^k$, we have

$$n - 2^{k-1} + 1 \geq \frac{n}{2} + 1.$$

This implies that in any Hamiltonian path, there exists an edge between two nodes with colour k . Then that edge must have colour k .

We next show that for each $1 \leq i < k$, every Hamiltonian path contains an edge of colour i . Suppose the contrary, that some Hamiltonian path does not contain an edge of colour i . Then nodes with colour i can only be adjacent to nodes with colour less than i inside the Hamiltonian path. Since there are 2^{i-1} nodes with colour i and $2^{i-1} - 1$ nodes with colour less than i , the Hamiltonian path must take the form

$$(i) \leftrightarrow (< i) \leftrightarrow (i) \leftrightarrow (< i) \leftrightarrow \dots \leftrightarrow (< i) \leftrightarrow (i),$$

where (i) denotes a node with colour i , $(< i)$ denotes a node with colour less than i , and \leftrightarrow denotes an edge. But this is impossible, as the Hamiltonian path would not have any nodes with colours greater than i . \square

Fix a set of k colours, we now prove that if there exists a good colouring of K_n , then $k \leq \lfloor \log_2 n \rfloor$. For $n = 2$, this is trivial, so we assume $n \geq 3$. For any node v of K_n and $1 \leq i \leq k$, we denote by $d_i(v)$ the number of edges with colour i incident with the node v .

Lemma 1. Consider a good colouring of K_n , and let AB be an arbitrary edge with colour i . If $d_i(A) + d_i(B) \leq n - 1$, then the colouring will remain good after recolouring edge AB with any other colour.

Proof. Suppose there exists a good colouring together with an edge AB of colour i , such that if AB is recoloured with another colour, the colouring will no longer be good. The failure of the new colouring being good will come from colour i , and thus there exists a Hamiltonian path containing edge AB such that initially (i.e. before recolouring) AB is the only edge of colour i in this path. Writing $A = A_0$ and $B = B_0$, denote this Hamiltonian path by

$$A_s \leftrightarrow A_{s-1} \leftrightarrow \dots \leftrightarrow A_1 \leftrightarrow A_0 \leftrightarrow B_0 \leftrightarrow B_1 \leftrightarrow \dots \leftrightarrow B_{t-1} \leftrightarrow B_t,$$

where $s, t \geq 0$ and $s + t + 2 = n$.

In the initial colouring, we observe the following.

- The edge B_0A_s must have colour i , since otherwise the path

$$A_0 \leftrightarrow A_1 \leftrightarrow \cdots \leftrightarrow A_{s-1} \leftrightarrow A_s \leftrightarrow B_0 \leftrightarrow B_1 \leftrightarrow \cdots \leftrightarrow B_{t-1} \leftrightarrow B_t$$

has no edges of colour i .

- Similarly, the edge A_0B_t must have colour i .
- For each $0 \leq p < s$, at least one of the edges B_0A_p and A_0A_{p+1} must have colour i , since otherwise the path

$$A_s \leftrightarrow \cdots \leftrightarrow A_{p+2} \leftrightarrow A_{p+1} \leftrightarrow A_0 \leftrightarrow A_1 \leftrightarrow \cdots \leftrightarrow A_{p-1} \leftrightarrow A_p \leftrightarrow B_0 \leftrightarrow B_1 \leftrightarrow \cdots \leftrightarrow B_t$$

has no edges of colour i .

- Similarly, for each $0 \leq q < t$, at least one of the edges A_0B_q and B_0B_{q+1} must have colour i .

In the above list, each edge A_0X appears exactly once and also each edge B_0X appears exactly once (where A_0B_0 and B_0A_0 are counted separately). Adding up the contributions to $d_i(A) + d_i(B)$, we obtain

$$d_i(A) + d_i(B) \geq (s + 1) + (t + 1) = n.$$

This contradicts our assumption that $d_i(A) + d_i(B) \leq n - 1$. □

Our strategy now is to repeatedly recolour the edges using Lemma 1 until the colouring has a simple structure. For a node v , we define $m(v)$ to be the largest value of $d_i(v)$ over all colours i .

Lemma 2. Assume we have a good colouring of K_n . Let A, B be two distinct nodes, and let j be the colour of edge AB where $1 \leq j \leq k$. If

- $m(A) \geq m(B)$ and
- $m(A) = d_i(A)$ for some $i \neq j$,

then after recolouring edge AB with colour i , the colouring remains good.

Proof. Note that

$$d_j(A) + d_j(B) \leq (n - 1 - m(A)) + m(B) \leq n - 1,$$

and so we may apply Lemma 1. □

Lemma 3. Assume we have a good colouring of K_n . Let S be a nonempty set of nodes. Let $A \in S$ be a node such that $m(A) \geq m(B)$ for all $B \in S$, and choose $1 \leq i \leq k$ for which $d_i(A) = m(A)$. Then after recolouring the edge AB with colour i for all $B \in S$ distinct from A , the colouring remains good.

Proof. We repeatedly perform the following operation until all edges AB with $B \in S$ have colour i :

choose an edge AB with $B \in S$ that does not have colour i , and recolour it with colour i .

By Lemma 2, the colouring remains good after one operation. Moreover, $m(A)$ increase by 1 during an operation, and all other $m(B)$ may increase by at most 1. This shows that $m(A)$ will remain maximal amongst $m(B)$ for $B \in S$. We will also have $d_i(A) = m(A)$ after the operation, since both sides increase by 1. Therefore the operation can be performed repeatedly, and the colouring remains good. □

We first apply Lemma 3 to the set of all n nodes in K_n . After recolouring, there exists a node A_1 such that every edge incident with A_1 has colour c_1 . We then apply Lemma 3 to the set of nodes excluding A_1 , and we obtain a colouring where

- every edge incident with A_1 has colour c_1 ,
- every edge incident with A_2 except for A_1A_2 has colour c_2 .

Repeating this process, we arrive at the following configuration:

- the n nodes of K_n are labelled A_1, A_2, \dots, A_n ,
- the node A_i has a corresponding colour c_i (as a convention, we also colour A_i with c_i),
- for all $1 \leq u < v \leq n$, the edge between A_u and A_v has colour c_u ,
- this colouring is good.

Claim 2. For every colour i , there exists a $1 \leq p \leq n$ such that the number of nodes of colour i amongst A_1, \dots, A_p is greater than $p/2$.

Proof. Suppose the contrary, that for every $1 \leq p \leq n$, there are at most $\lfloor p/2 \rfloor$ nodes of colour i . We then construct a Hamiltonian path not containing any edge of colour i . Let A_{x_1}, \dots, A_{x_t} be the nodes with colour i , where $x_1 < x_2 < \dots < x_t$, and let $A_{y_1}, A_{y_2}, \dots, A_{y_s}$ be the nodes with colour different from i , where $y_1 < y_2 < \dots < y_s$. We have $s + t = n$ and $t \leq \lfloor n/2 \rfloor$, so $t \leq s$. We also see that $y_j < x_j$ for all $1 \leq j \leq t$, because otherwise, A_1, A_2, \dots, A_{x_j} will have j nodes of colour i and less than j nodes of colour different from i . Then we can construct a Hamiltonian path

$$A_{x_1} \leftrightarrow A_{y_1} \leftrightarrow A_{x_2} \leftrightarrow A_{y_2} \leftrightarrow A_{x_3} \leftrightarrow \dots \leftrightarrow A_{x_t} \leftrightarrow A_{y_t} \leftrightarrow A_{y_{t+1}} \leftrightarrow \dots \leftrightarrow A_{y_s}$$

that does not contain an edge with colour i . This contradicts that the colouring is good. \square

So for every colour i , there has to be an integer p_i with $1 \leq p_i \leq n$ such that there are more than $p_i/2$ nodes assigned colour i amongst A_1, \dots, A_{p_i} . Choose the smallest such p_i for every i , and without loss of generality assume

$$p_1 < p_2 < \dots < p_k.$$

Note that the inequalities are strict by the definition of p_i .

Then amongst the nodes A_1, \dots, A_{p_i} , there are at least $\lceil (p_i + 1)/2 \rceil$ nodes of colour j for all $1 \leq j \leq i$. Then

$$p_i \geq \left\lceil \frac{p_1 + 1}{2} \right\rceil + \left\lceil \frac{p_2 + 1}{2} \right\rceil + \dots + \left\lceil \frac{p_i + 1}{2} \right\rceil.$$

This inductively shows that

$$p_i \geq 2^i - 1$$

for all $1 \leq i \leq k$, and this already proves $n \geq 2^k - 1$.

It remains to show that $n = 2^k - 1$ is not possible. If $n = 2^k - 1$, then all inequalities have to be equalities, so $p_i = 2^i - 1$ and there must be exactly 2^{i-1} nodes of colour i . Moreover, there cannot be a node of colour i amongst $A_1, A_2, \dots, A_{p_{i-1}}$, and so the set of nodes of colour i must precisely be

$$A_{2^{i-1}}, A_{2^{i-1}+1}, \dots, A_{2^i-1}.$$

Then we can form a Hamiltonian path

$$A_{2^k-1} \leftrightarrow A_1 \leftrightarrow A_{2^{k-1}+1} \leftrightarrow A_2 \leftrightarrow A_{2^{k-1}+2} \leftrightarrow A_3 \leftrightarrow \dots \leftrightarrow A_n,$$

which does not contain an edge of colour k . This is a contradiction, and therefore $n \geq 2^k$.


Geometry

G1. Let $ABCDE$ be a convex pentagon such that $\angle ABC = \angle AED = 90^\circ$. Suppose that the midpoint of CD is the circumcentre of triangle ABE . Let O be the circumcentre of triangle ACD .

Prove that line AO passes through the midpoint of segment BE .

(Slovakia)

Solution 1 (Area Ratio).



Let M be the midpoint of CD and $X = BC \cap ED$. Since $\angle ABX = \angle AEX = 90^\circ$, AX is a diameter of the circumcircle of $\triangle ABE$ so the midpoint of AX is the circumcentre of $\triangle ABE$. Therefore, the midpoint of AX coincides with M . This means $ACXD$ is a parallelogram and in particular, $AD \parallel BC$ and $AC \parallel ED$.

We denote the area of $\triangle P_1P_2P_3$ by $[P_1P_2P_3]$. To prove that line AO bisects BE , it suffices to show $[OAB] = [OAE]$.

Let C', D' be the midpoints of AC, AD respectively. Since $OD' \perp AD$, $AD \parallel BC$, and $BC \perp AB$, we have $AB \parallel OD'$, so $[OAB] = [D'AB]$. Using $AD \parallel BC$ again, we have $[D'AB] = [D'AC]$. Therefore

$$[OAB] = [D'AB] = [D'AC] = \frac{1}{2}[ACD].$$

Similarly


$$[OAE] = [C'AE] = [C'AD] = \frac{1}{2}[ACD].$$

Combining these gives $[OAB] = [OAE]$.

Comment 1. The following is another way to prove $AD \parallel BC$ and $AC \parallel ED$.

Let ℓ be the perpendicular bisector of AB . Since the midpoint of CD is the circumcentre of $\triangle ABE$, it must lie on ℓ . Also, since $\angle ABC = 90^\circ$, the midpoint of AC is on ℓ . Therefore, we get $AD \parallel \ell \parallel BC$ and similarly $AC \parallel ED$.

Solution 2 (Similar Triangles).



Let M be the midpoint of CD and $X = BC \cap ED$. As in Solution 1, M is the midpoint of AX and $ACXD$ is a parallelogram. Since $AD \parallel BC$ and $\angle ABC = 90^\circ$, we have $\angle DAB = 90^\circ$.


Let N be the midpoint of BE . It is enough to show that $\angle NAB = \angle OAB$. Since $ABXE$ is cyclic, we have

$$\angle ABE = \angle AXE = \angle XAC \quad \text{and} \quad \angle BEA = \angle CXA.$$

Therefore, $\triangle ABE \sim \triangle CAX$, and N corresponds to M under this similarity. In particular, $\angle NAB = \angle ACM$.

Also, we have

$$\angle OAB = 90^\circ - \angle DAO = \angle ACM = \angle NAB.$$

Solution 3 (Reflection).

Let N be the midpoint of BE , and let F, G be the projections of C, D onto AD, AC respectively. $CGFD$ is cyclic so

$$\angle AGF = \angle CDF = \angle CDA = 90^\circ - \angle OAC$$

giving $AO \perp FG$. Therefore it's enough to show that $AN \perp FG$.

As in Solution 1, $AD \parallel BC$ and $AC \parallel ED$ so $\angle EAG = \angle FAB = 90^\circ$ and in fact $AEDG$ and $AFCB$ are rectangles. From this we get

$$\angle AGE = \angle DAG = \angle FAC = \angle BFA$$

so $\triangle GAE \sim \triangle FAB$.

Let F' be the reflection of F in A , then $\triangle F'AB \sim \triangle FAB \sim \triangle GAE$. Thus A is the centre of the spiral symmetry taking $F'B \rightarrow GE$.

Let P be the midpoint of $F'G$ then by the spiral similarity, we have $\triangle APN \sim \triangle AGE$ which implies $\angle NAP = 90^\circ$. From A being the midpoint of FF' we have $AP \parallel FG$. Combining the results gives $AN \perp FG$.


G2.

Let ABC be a triangle with $AC > BC$. Let ω be the circumcircle of triangle ABC and let r be the radius of ω . Point P lies on segment AC such that $BC = CP$ and point S is the foot of the perpendicular from P to line AB . Let ray BP intersect ω again at D and let Q lie on line SP such that $PQ = r$ and S, P, Q lie on the line in that order. Finally, let the line perpendicular to CQ from A intersect the line perpendicular to DQ from B at E .

Prove that E lies on ω .

(Iran)

Solution 1 (Similar Triangles).



First observe that

$$\angle DPA = \angle BPC \stackrel{CP=CB}{=} \angle CBP = \angle CBD = \angle CAD = \angle PAD$$

so $DP = DA$. Thus there is a symmetry in the problem statement swapping $(A, D) \leftrightarrow (B, C)$.

Let O be the centre of ω and let E be the reflection of P in CD which, by

$$\angle CED = \angle DPC = 180^\circ - \angle CPB \stackrel{CP=CB}{=} 180^\circ - \angle PBC = 180^\circ - \angle DBC$$


lies on ω . We claim the two lines concur at E . By the symmetry noted above, it suffices to prove that $BE \perp DQ$ and then $AE \perp CQ$ will follow by symmetry.

We have $AO = PQ$, $AD = DP$ and

$$\angle DAO = 90^\circ - \angle ABD \stackrel{PQ \perp AB}{=} \angle DPQ.$$

Hence $\triangle AOD \cong \triangle PQD$. Thus

$\angle QDB + \angle DBE = \angle ODA + \angle DAE \stackrel{DE=DA}{=} \angle ODA + \angle AED = (90^\circ - \angle AED) + \angle AED = 90^\circ$
giving $BE \perp DQ$ as required.

Solution 2 (Second Circle).

As in Solution 1, we prove that $DA = DP$ and note the symmetry in the problem statement swapping $(A, D) \leftrightarrow (B, C)$.

Let Γ be the circumcircle of $\triangle PCD$. Since $DP = DA$ and $\angle ACD = \angle PCD$, the radius of Γ is equal to that of ω . We have that

$$\angle DPQ = \angle BPS = 90^\circ - \angle ABD = 90^\circ - \angle PCD.$$

This, combined with PQ being equal to the common circumradius of Γ and ω , means that Q is the circumcentre of Γ .

Let the perpendiculars to CQ, DQ from A, B intersect at E then we have

$$\begin{aligned} \angle EAC &= 90^\circ - \angle ACQ \stackrel{QC=QP}{=} 90^\circ - \angle QPC = 90^\circ - \angle SPA = \angle CAB \implies \angle EAB = 2\angle PAB \\ \angle DBE &= 90^\circ - \angle QDP \stackrel{QD=QP}{=} 90^\circ - \angle DPQ = 90^\circ - \angle BPS = \angle ABD \implies \angle ABE = 2\angle ABP. \end{aligned}$$

Combining these

$$\angle BEA = 180^\circ - 2(\angle PAB + \angle ABP) = 180^\circ - 2\angle APD \stackrel{DA=DP}{=} \angle BDA$$

which gives that E lies on ω .

Comment 1. An alternative final angle chase is

$$\angle BEA = 180^\circ - \angle CQD \stackrel{\Gamma}{=} 180^\circ - 2(180^\circ - \angle DPC) = 180^\circ - 2\angle APD \stackrel{DA=DP}{=} \angle PDA = \angle BDA.$$

Comment 2. An alternative formulation of the problem in terms of a cyclic quadrilateral is given below:

Let $ABCD$ be a cyclic quadrilateral with circumcircle ω and circumradius r . The diagonals AC and BD intersect at P . Suppose that $AD = DP$. Let S be the foot of the perpendicular from P to the line AB . Point Q lies on line SP such that $PQ = r$ and S, P, Q lie on the line in that order. Let the line perpendicular to CQ from A intersect the line perpendicular to DQ from B at E .

Prove that E lies on ω .

G3. Let $ABCD$ be a cyclic quadrilateral with $\angle BAD < \angle ADC$. Let M be the midpoint of the arc CD not containing A . Suppose there is a point P inside $ABCD$ such that $\angle ADB = \angle CPD$ and $\angle ADP = \angle PCB$.


Prove that lines AD, PM, BC are concurrent.

(Slovakia)

Solution 1. Let X and Y be the intersection points of AM and BM with PD and PC respectively. Since $ABCD$ is cyclic and $CM = MD$, we have


$$\angle XAD = \angle MAD = \angle CBM = \angle CBY.$$

Combining this with $\angle ADX = \angle YCB$, we get $\angle DXA = \angle BYC$, and so $\angle PXM = \angle MYP$. Moreover, $\angle YPX = \angle CPD = \angle ADB = \angle AMB$. The quadrilateral $MXPY$ therefore has equal opposite angles and so is a parallelogram.



Let R and S be the intersection points of AM and BM with BC and AD respectively. Due to $AM \parallel PC$ and $BM \parallel PD$, we have $\angle ASB = \angle ADP = \angle PCB = \angle ARB$ and so the quadrilateral $ABRS$ is cyclic. We then have $\angle SRB = 180^\circ - \angle BAS = \angle DCB$ and so $SR \parallel CD$. In triangles PCD and MRS , the corresponding sides are parallel so they are homothetic meaning lines DS, PM, CR concur at the centre of this homothety.

Solution 2. Let AD and BC meet at T . Denote by p_a, p_b, m_a and m_b the distances between line TA and P , TB and P , TA and M and between TB and M respectively. Our goal is to prove $p_a : p_b = m_a : m_b$ which is equivalent to the collinearity of T, P and M .



Let $\angle BAC = \angle BDC = \alpha$, $\angle DBA = \angle DCA = \beta$, $\angle ADB = \angle AMB = \angle ACB = \angle CPD = \mu$, $\angle ADP = \angle PCB = \nu$ and $\angle MAD = \angle CAM = \angle MBD = \angle CBM = \chi$.

From $\angle ADP = \angle PCB = \nu$ and $\angle MAD = \angle CBM = \chi$ we get

$$\frac{p_a}{p_b} = \frac{PD \cdot \sin \nu}{PC \cdot \sin \nu} = \frac{PD}{PC} \quad \text{and} \quad \frac{m_a}{m_b} = \frac{MA \cdot \sin \chi}{MB \cdot \sin \chi} = \frac{MA}{MB}.$$

Hence $p_a : p_b = m_a : m_b$ is equivalent to $PD : PC = MA : MB$, and since $\angle CPD = \angle AMB = \mu$, this means we have to show that triangles PDC and MAB are similar.

In triangle PDC we have

$$\begin{aligned} \angle PDC + \angle DCP &= 180^\circ - \angle CPD = 180^\circ - \mu, \\ \angle PDC - \angle DCP &= (\alpha + \mu - \nu) - (\beta + \mu - \nu) = \alpha - \beta. \end{aligned}$$

Similarly, in triangle MAB we have

$$\begin{aligned} \angle BAM + \angle MBA &= 180^\circ - \angle AMB = 180^\circ - \mu, \\ \angle BAM - \angle MBA &= (\alpha + \chi) - (\beta + \chi) = \alpha - \beta. \end{aligned}$$

Therefore, $(\angle BAM, \angle MBA)$ and $(\angle PDC, \angle DCP)$ satisfy the same system of linear equations. The common solution is

$$\angle BAM = \angle PDC = \frac{180^\circ - \mu + \alpha - \beta}{2} \quad \text{and} \quad \angle MBA = \angle DCP = \frac{180^\circ - \mu - \alpha + \beta}{2}.$$

Hence triangles PDC and MAB have equal angles and so are similar. This completes the proof.


G4. Let ABC be an acute-angled triangle with $AB < AC$. Denote its circumcircle by Ω and denote the midpoint of arc CAB by S . Let the perpendicular from A to BC meet BS and Ω at D and $E \neq A$ respectively. Let the line through D parallel to BC meet line BE at L and denote the circumcircle of triangle BDL by ω . Let ω meet Ω again at $P \neq B$.

Prove that the line tangent to ω at P , and line BS intersect on the internal bisector of $\angle BAC$.

(Portugal)

Solution 1 (Triangles in Perspective). Let S' be the midpoint of arc BC of Ω , diametrically opposite to S so SS' is a diameter in Ω and AS' is the angle bisector of $\angle BAC$. Let the tangent of ω at P meet Ω again at $Q \neq P$, then we have $\angle S QS' = 90^\circ$.

We will show that triangles APD and $S'QS$ are similar and their corresponding sides are parallel. Then it will follow that the lines connecting the corresponding vertices, namely line AS' , that is the angle bisector of $\angle BAC$, line PQ , that is the tangent to ω at P , and DS are concurrent. Note that the sides AD and $S'S$ have opposite directions, so the three lines cannot be parallel.



First we show that $AP \perp DP$. Indeed, from cyclic quadrilaterals $APBE$ and $DPLB$ we can see that

$$\angle PAD = \angle PAE = 180^\circ - \angle EBP = \angle PBL = \angle PDL = 90^\circ - \angle ADP.$$

Then, in triangle APD we have $\angle DPA = 180^\circ - \angle PAD - \angle ADP = 90^\circ$.

Now we can see that:

- Both lines ADE and SS' are perpendicular to BC , so $AD \parallel S'S$.
- Line PQ is tangent to circle ω at P so $\angle DPQ = \angle DBP = \angle SBP = \angle SQP$; it follows that $PD \parallel QS$.


- Finally, since $AP \perp PD \parallel QS \perp S'Q$, we have $AP \parallel S'Q$ as well.

Hence the corresponding sides of triangles APD and $S'QS$ are parallel completing the solution.

Solution 2 (Pascal). Again, let S' be the midpoint of arc BC , diametrically opposite to S , so $AES'S$ is an isosceles trapezoid, and $\angle S'BS = \angle S'PS = 90^\circ$. Let lines AE and PS' meet at T and let AP and $S'B$ meet at point M .

We will need that points L, P, S are collinear, and points T and M lie on circle ω .

- From $\angle LPB = \angle LDB = 90^\circ - \angle BDE = 90^\circ - \angle BSS' = \angle SS'B = 180^\circ - \angle BPS$ we get $\angle LPB + \angle BPS = 180^\circ$, so L, P and S are indeed collinear.
- Since SS' is a diameter in Ω , lines LPS and PTS' are perpendicular. We also have $LD \parallel BC \perp AE$ hence $\angle LDT = \angle LPT = 90^\circ$ and therefore $T \in \omega$.
- By $\angle LPM = \angle SPA = \angle SEA = \angle EAS' = \angle EBS' = \angle LBM$, point M is concyclic with B, P, L so $M \in \omega$.



Now let X be the intersection of line BDS with the tangent of ω at P and apply Pascal's theorem to the degenerate cyclic hexagon $PPMBDT$. This gives points $PP \cap BD = X$, $PM \cap DT = A$ and $MB \cap TP = S'$ are collinear so X lies on line AS' , that is the bisector of $\angle BAC$.


Comment. It is easy to see that $LMTD$ is a rectangle, but we did not need this information for the solution.

Solution 3 (Projections). Let A' and S' be the points of Ω diametrically opposite to A and S respectively. It is well-known that E and A' are reflections with respect to SS' so AS' is the angle bisector of $\angle EAA'$. Define point T to be the intersection of AE and PS' . As in the previous two solutions, we have: $\angle DPA = 90^\circ$ so PD passes through A' ; points L, P, S are collinear; and $T \in \omega$.

Let lines AS' and PDA' meet at R . From the angles of triangles PRS' and PTE we get

$$\angle ARP = \angle AS'P + \angle S'PA' = \angle AEP + \angle EPS' = \angle ATP$$

so points A, P, T, R are concyclic. Denote their circle by γ . Due to $\angle RPA = \angle DPA = 90^\circ$, segment AR is a diameter in γ .



We claim that circles ω and γ are perpendicular. Let line LPS meet γ again at $U \neq P$, and consider triangles PLT and PTU . By $\angle LPT = \angle TPU = 90^\circ$ and

$$\angle PTL = \angle PBL = 180^\circ - \angle EBP = \angle PAE = \angle PAT = \angle PUT,$$

triangles PLT and PTU are similar. It follows that the spiral similarity that takes PLT to PTU , maps ω to γ and the angle of this similarity is 90° , so circles ω and γ are indeed perpendicular.

Finally, let lines BDS and ARS' meet at X . We claim that X bisects AR , so point X is the centre of γ and, as ω and γ are perpendicular, PX is tangent to ω .

Let t be the tangent of ω at D . From $\angle(DT, t) = \angle TPD = \angle S'PA' = \angle EAS'$ it can be seen that $t \parallel AS'$. Let I be the common point at infinity of t and AS' . Moreover, let lines LPS and $ADTE$ meet at V . By projecting line AS' to circle ω through D , then projecting ω to line AE through L , finally projecting AE to Ω through P , we find

$$\frac{AX}{RX} = (A, R; X, I) \stackrel{D}{=} (T, P; B, D) \stackrel{L}{=} (T, V; E, D) \stackrel{P}{=} (S', S; E, A') = -1,$$

so X is the midpoint of AR .

G5. Let ABC be an acute-angled triangle with circumcircle ω and circumcentre O . Points $D \neq B$ and $E \neq C$ lie on ω such that $BD \perp AC$ and $CE \perp AB$. Let CO meet AB at X , and BO meet AC at Y .

Prove that the circumcircles of triangles BXD and CYE have an intersection on line AO .
(Malaysia)

Solution 1 (Reflections).

Note that $AO = OC$ implies the lines AO , XO are reflections of each other about the line parallel to AC through O , which is the perpendicular bisector of BD . Call this line ℓ .


Let $P \neq X$ be the second intersection of circle $\odot BXD$ with line XO , and let Z be the intersection of circle $\odot BXD$ with line AO furthest from A .


Consider a reflection across ℓ . This maps B to D , AO to XO , and circle $\odot BXD$ to itself so the transformation must map P , the intersection of XO and circle $\odot BXD$, to the intersection of AO and $\odot BXD$ furthest from A i.e. Z . Thus we have

$$\angle OZB = \angle DPO = \angle DPX = \angle DBX = 90^\circ - \angle BAC = \angle OCB$$

which implies $BOCZ$ is cyclic.

Therefore the second intersection of circle $\odot BOC$ with line AO lies on circle $\odot BXD$. Similarly, Z lies on circle $\odot CYE$ so the two circles have common point Z on AO .



Solution 2 (Similar Triangles).

Let B' be the reflection of B in AC and let AO intersect circle $\odot OBC$ again at $Z \neq O$. Observe that

$$\angle B'CA + \angle ACZ = 2\angle ACB + \angle BCZ = 2\angle ACB + \angle BOZ = 2\angle ACB + (180^\circ - \angle AOB) = 180^\circ$$

so Z, C, B' are collinear.

Claim. Triangles ZXA and ZDB' are similar.

Proof. We have

$$\angle XAZ = \angle BAO = 90^\circ - \angle ACB = \angle CBB' = \angle BB'C = \angle DB'Z.$$

So it suffices to prove that $\frac{B'Z}{B'D} = \frac{AZ}{AX}$. To do this, first observe

$$\angle B'ZA = \angle CZO = \angle CBO = \angle XCB \quad \text{and} \quad \angle AB'Z = \angle AB'C = \angle CBA = \angle CBX.$$

Hence triangles ZAB' and CXB are similar so

$$\frac{B'Z}{AZ} = \frac{BC}{CX}.$$

Note that the orthocentre H of triangle ABC is the reflection of D in AC . Applying sine rule to triangles ACX and BHC gives

$$\frac{AX}{CX} = \frac{\sin \angle ACX}{\sin \angle XAC} = \frac{\sin (90^\circ - \angle CBA)}{\sin \angle BAC} = \frac{\sin (90^\circ - \angle CBA)}{\sin (180^\circ - \angle BAC)} = \frac{\sin \angle HCB}{\sin \angle BHC} = \frac{BH}{BC} = \frac{B'D}{BC}.$$

Multiplying the two results gives

$$\frac{B'Z}{AZ} \cdot \frac{AX}{CX} = \frac{BC}{CX} \cdot \frac{B'D}{BC} = \frac{B'D}{CX}$$


which implies $\frac{B'Z}{B'D} = \frac{AZ}{AX}$, as required. \square

From the claim

$$\angle BDZ = 180^\circ - \angle ZDB' = 180^\circ - \angle ZXA = \angle BXZ$$

which means Z lies on circle $\odot BXD$. Similarly, Z lies on circle $\odot CYE$ completing the proof.

Solution 3 (Inversion at A).



Consider the composition of the inversion at A with radius $\sqrt{AB \times AC}$ and reflection in the angle bisector of $\angle BAC$, and use P^* to denote the image of a point P under this transformation. Let H be the orthocentre of triangle ABC and let K, L be the feet of the perpendicular from A, B to BC, CA respectively. Denote $A = \angle BAC$, $B = \angle CBA$ and $C = \angle ACB$.

We have

$$\angle D^*AK = \angle OAD = 90^\circ - \angle DBA = 90^\circ - (90^\circ - A) = A.$$

Hence, using right-angled $\triangle AKD^*$

$$\begin{cases} D^*K = AK \tan A = 2R \sin B \sin C \tan A \\ HK = 2R \cos B \cos C \end{cases} \implies \frac{D^*K}{HK} = \tan A \tan B \tan C$$

We also have

$$\angle AX^*B = \angle ACX = \angle ACO = 90^\circ - B.$$

Hence, using right-angled $\triangle BLX^*$

$$\begin{cases} X^*L = BL \tan B = 2R \sin A \sin C \tan B \\ HL = 2R \cos A \cos C \end{cases} \implies \frac{X^*L}{HL} = \tan A \tan B \tan C.$$

Thus $\frac{D^*K}{HK} = \frac{X^*L}{HL}$ and as $\angle HLX^* = \angle HKD^* = 90^\circ$, this means that triangle D^*HK and X^*HL are similar and in particular

$$\angle CD^*H = \angle KD^*H = \angle LX^*H = \angle CX^*H$$


so D^*HCX^* is cyclic.

Inverting back, this gives DH^*BX cyclic so H^* lies on circle $\odot BXD$. Similarly, H^* lies on circle $\odot CYE$.

Since AO and AH are isogonal in $\angle BAC$, H^* lies on line AO completing the proof.

Solution 4 (Inversion at O). Let F be the point on ω such that AF is a diameter of ω , and J be the intersection of DF with CO .

Consider the inversion with respect to ω and use P' to denote the image of a point P .



X' lies on line CO and we have

$$\angle BX'J = \angle BX'O = \angle OBX = \angle OBA = \angle BAO = \angle BAF = \angle BDF = \angle BDJ$$

so $BX'DJ$ is cyclic.

Let K be the intersection of AF with BC . Then we have $OB = OD$ and

$$\begin{aligned} \angle KBO &= 90^\circ - A = \angle DBA = \angle DFA = \angle DFO = \angle ODJ \\ \angle BOK &= 2\angle OBA = 2\angle CBD = \angle COD = \angle JOD \end{aligned}$$

Hence triangle BOK and DOJ are congruent. In particular $BK = DJ$ and

$$\angle KBD = \angle KBO + \angle OBD = \angle ODJ + \angle BDO = \angle BDJ.$$

Thus $BDJK$ is an isosceles trapezoid and $BX'DJK$ is cyclic.


Inverting back this gives that $BXDK'$ is cyclic. Similarly $CYEK'$ is cyclic. Since K lies on AO , K' also lies on AO completing the proof.

G6. Let ABC be an acute-angled triangle with circumcircle ω . A circle Γ is internally tangent to ω at A and also tangent to BC at D . Let AB and AC intersect Γ at P and Q respectively. Let M and N be points on line BC such that B is the midpoint of DM and C is the midpoint of DN . Lines MP and NQ meet at K and intersect Γ again at I and J respectively. The ray KA meets the circumcircle of triangle IJK at $X \neq K$.

Prove that $\angle BXP = \angle CXQ$.

(United Kingdom)

Solution 1 (Similar Triangles).



Let MP and NQ intersect AD at K_1 and K_2 respectively. By applying Menelaus' theorem to triangle ABD and line MPK_1 , we have

$$\frac{AK_1}{K_1D} = \frac{AP}{PB} \cdot \frac{BM}{MD} = \frac{AP}{2PB}$$

and similarly $\frac{AK_2}{K_2D} = \frac{AQ}{2QC}$. A homothety at A takes $\Gamma \rightarrow \omega$ and D to the midpoint of arc BC not containing A , so $PQ \parallel BC$ and AD bisects $\angle BAC$. Thus

$$\frac{AK_1}{K_1D} = \frac{AP}{2PB} = \frac{AQ}{2QC} = \frac{AK_2}{K_2D}$$

which implies $K_1 \equiv K_2$, and K lies on AD .

Then we obtain

$$\angle JXD = \angle JXK = \angle JIK = \angle JIP = \angle JQP = \angle JND$$

where the last equality follows from $PQ \parallel BC$. This shows $JXND$ is cyclic and hence

$$\angle DXN = \angle DJN = \angle DJQ = \angle DAQ = \angle DAC$$

which shows $AC \parallel XN$. As C is the midpoint of DN , A is the midpoint of XD .

Now observe that $\angle ADP = \angle AQP = \angle ACB$ and $\angle PAD = \angle DAC = \frac{\angle A}{2}$ so triangle APD and ADC are similar. Therefore we have

$$\frac{CD}{DP} = \frac{AD}{AP} = \frac{XA}{AP}$$

and also have

$$\angle CDP = 180^\circ - \angle PDB = 180^\circ - \angle PAD = \angle XAP.$$

Combining the two results gives triangles PDC and PAX are similar, which shows P is the centre of spiral similarity taking $CD \rightarrow XA$. Hence also triangles PXC and PAD are similar which shows $\angle PXC = \angle PAD = \frac{\angle A}{2}$. This gives


$$\angle BXP = \angle BXC - \angle PXC = \angle BXC - \frac{\angle A}{2}$$

which is symmetric in B, C giving the result.

Solution 2 (Inversion). As in the first solution we show that K lies on AD . From C being the midpoint of DN and $BC \parallel PQ$ we get

$$-1 = (C, \infty_{BC}; N, D) \stackrel{Q}{=} (A, PQ \cap AD; K, D) \stackrel{Q}{=} (A, P; J, D).$$

Similarly we get $(A, Q; I, D) = -1$.



Now invert about D with radius $DP = DQ$ denoting the inverse of a point Z by Z^* . Since $\odot APQ$ and line PQ swap we have $A^* = PQ \cap AD$. Thus we have:

$$-1 = (A, A^*; K, D) = (A, P; J, D) = (A, Q; I, D)$$

As inversion preserves cross ratio and D inverts to the point at infinity, it follows I^*, J^*, K^* are the midpoints of A^*Q, A^*P, A^*A respectively. We know $XIKJ$ cyclic so X is the second intersection of circle $(I^*J^*K^*)$ with AD . Homothety of factor 2 at A^* takes circle $(I^*J^*K^*)$ to circle $\odot APQ \equiv \Gamma$ hence in fact X^* is the midpoint of A^*D .

Then we have

$$\angle PDB = \angle BAD = \angle DAC = \angle DC^*A^*$$

so $DP \parallel C^*A^*$. Also A^* lies on PQ so as $PQ \parallel BC$ we get $A^*P \parallel DC^*$, which gives PA^*C^*D is a parallelogram. Similarly QA^*B^*D is also a parallelogram. As X^* is the midpoint of A^*D this shows that X^* lies on lines B^*Q and C^*P .

By applying standard properties of angles under inversion, we have

$$\begin{aligned} \angle BXP - \angle CXQ &= (\angle BXD - \angle PXD) - (\angle DXC - \angle DXQ) \\ &= (\angle DB^*X^* - \angle DPX^*) - (\angle X^*C^*D - \angle X^*QD) \\ &= (\angle DB^*X^* + \angle X^*PQ) - (\angle X^*C^*D + \angle PQX^*) \\ &\quad \text{(as } \angle DPQ = \angle PQD) \\ &= \underbrace{(\angle DB^*Q - \angle PQB^*)}_{=0} - \underbrace{(\angle PC^*D - \angle C^*PQ)}_{=0} \\ &= 0 \quad \text{(as } PQ \parallel BC) \end{aligned}$$


which gives the result.

G7. Let ABC be an acute, scalene triangle with orthocentre H . Let ℓ_a be the line through the reflection of B with respect to CH and the reflection of C with respect to BH . Lines ℓ_b and ℓ_c are defined similarly. Suppose lines ℓ_a , ℓ_b , and ℓ_c determine a triangle \mathcal{T} .

Prove that the orthocentre of \mathcal{T} , the circumcentre of \mathcal{T} and H are collinear.

(Ukraine)

Solution 1.



We write $\triangle P_1P_2P_3 \stackrel{+}{\sim} \triangle Q_1Q_2Q_3$ (resp. $\triangle P_1P_2P_3 \stackrel{-}{\sim} \triangle Q_1Q_2Q_3$) to indicate that two triangles are directly (resp. oppositely) similar. We use directed angles throughout denoted with \sphericalangle .

Denote by A_b , A_c the reflections of A in BH and CH respectively. B_c , B_a and C_a , C_b are defined similarly. By definition, $\ell_a = B_cC_b$, $\ell_b = C_aA_c$, $\ell_c = A_bB_a$. Let $A_1 = \ell_b \cap \ell_c$, $B_1 = \ell_c \cap \ell_a$, $C_1 = \ell_a \cap \ell_b$ and let O_1 , H_1 be the orthocentre and circumcentre of $\mathcal{T} \equiv \triangle A_1B_1C_1$ respectively.

Claim 1. $\triangle AA_bA_c \stackrel{-}{\sim} \triangle ABC$.

Proof. Let $P = BH \cap AC$, $Q = CH \cap AB$, then it is well known that $\triangle APQ \stackrel{-}{\sim} \triangle ABC$. By the dilation with factor 2 centred at A , $\triangle APQ$ is sent to $\triangle AA_bA_c$, so we have $\triangle AA_bA_c \stackrel{-}{\sim} \triangle ABC$. \square

Claim 2. $\triangle AA_bA_c \stackrel{+}{\sim} \triangle AB_aC_a$ and A_1 lies on the circumcircle of $\triangle AA_bA_c$ which is centred at H .

Proof. Since B_a, C_a are reflections of B, C in AH , we have $\triangle AB_aC_a \stackrel{\sim}{\sim} \triangle ABC$. Combining this with *Claim 1*, we have $\triangle AA_bA_c \stackrel{+}{\sim} \triangle AB_aC_a$, where A is the centre of this similarity. Therefore, $\sphericalangle A_cA_1A_b = \sphericalangle A_cAA_b$ meaning A_1 lies on $\odot AA_bA_c$. By symmetry, $HA_b = HA = HA_c$, so H is centre of this circle. \square

Claim 3. $\triangle A_1B_1C_1 \stackrel{\sim}{\sim} \triangle ABC$.

Proof. From *Claim 2* we have

$$\sphericalangle C_1A_1B_1 = \sphericalangle A_cA_1A_b \stackrel{\text{Claim 2}}{=} \sphericalangle A_cAA_b = -\sphericalangle CAB$$

and similarly $\sphericalangle A_1B_1C_1 = -\sphericalangle ABC$, $\sphericalangle B_1C_1A_1 = -\sphericalangle BCA$, which imply $\triangle A_1B_1C_1 \stackrel{\sim}{\sim} \triangle ABC$. \square

Denote the ratio of similitude of $\triangle A_1B_1C_1$ and $\triangle ABC$ by λ ($= \frac{B_1C_1}{BC}$), then

$$\lambda = \frac{H_1A_1}{HA} = \frac{H_1B_1}{HB} = \frac{H_1C_1}{HC}.$$

Since $HA = HA_1$ and similarly $HB = HB_1$, $HC = HC_1$ from *Claim 2*, we get


$$\lambda = \frac{H_1A_1}{HA_1} = \frac{H_1B_1}{HB_1} = \frac{H_1C_1}{HC_1}.$$

Therefore, the circle $A_1B_1C_1$ is the Apollonian circle of the segment HH_1 with ratio λ so the line HH_1 passes through O_1 .

Solution 2. We use the same notation $A_b, A_c, B_c, B_a, C_a, C_b$ and A_1, B_1, C_1, O_1, H_1 as in Solution 1, and also show *Claim 1*, *Claim 2* and *Claim 3* in the same way.

Let O be the circumcentre of $\triangle ABC$ and A_2 be the reflection of A_1 in AH . As $\odot AA_bA_c$ is centred at H , A_2 also lies on this circle.

By *Claim 2*, $\angle B_aA_1C_a = \angle A_bAA_c = \angle B_aAC_a$, so A_1 lies on $\odot AB_aC_a$. Reflecting this in AH gives that A_2 lies on $\odot ABC$. We now have circles centred at O and H passing through A and A_2 so these points are symmetric with respect to OH . Define B_2 and C_2 similarly then $\triangle ABC$ and $\triangle A_2B_2C_2$ are symmetric with respect to OH and also $\odot ABC = \odot A_2B_2C_2$.



Claim 4. A_1A_2 , B_1B_2 and C_1C_2 have an intersection on $\odot ABC$ which we denote by T .

Proof. Let $T = A_1A_2 \cap B_1B_2$. Since $A_1A_2 \parallel BC$ and $B_1B_2 \parallel AC$, we have

$$\angle A_2TB_2 = \angle BCA = -\angle B_2C_2A_2 = \angle A_2C_2B_2.$$

So T lies on $\odot A_2B_2C_2 = \odot ABC$. Similarly the intersection of A_1A_2 and C_1C_2 lies on $\odot ABC$, so C_1C_2 also passes through T . \square

Claim 5. T also lies on $\odot A_1B_1C_1$ and T corresponds to T itself under the similarity $\triangle A_1B_1C_1 \sim \triangle ABC$.

Proof. We know $\triangle A_1B_1C_1 \sim \triangle ABC$ by *Claim 3*. We also have

$$\sphericalangle B_1TC_1 = \sphericalangle B_2TC_2 = \sphericalangle B_2A_2C_2 = -\sphericalangle BAC \stackrel{\text{Claim 3}}{=} \sphericalangle B_1A_1C_1,$$

so T lies on $\odot A_1B_1C_1$. The remaining part is concluded by the following angle chase:

$$\sphericalangle A_1B_1T = \sphericalangle A_1B_1B_2 \stackrel{B_1B_2 \parallel AC}{=} \sphericalangle A_1A_bA = \sphericalangle A_1A_2A = -\sphericalangle AA_2T = -\sphericalangle ABT. \quad \square$$

Claim 6. The circumradius of $\triangle A_1B_1C_1$ is equal to HO .

Proof. Two circles centred at H intersect ℓ_c at A_1, A_b and B_1, B_a , so A_1A_b and B_1B_a have the same midpoint and thus $A_1B_1 = A_bB_a$. Consider the spiral symmetry $\triangle AA_bA_c \stackrel{+}{\sim} \triangle AB_aC_a$. This takes H , the circumcentre of $\triangle AA_bA_c$, to the circumcentre of $\triangle AB_aC_a$, denoted by O_a , which is symmetric to O in AH . Hence $\triangle AA_bB_a \stackrel{+}{\sim} \triangle AHO_a$, so

$$\frac{AA_b}{B_aA_b} = \frac{AH}{HO_a} = \frac{AH}{HO} \implies \frac{AH}{AA_b} = \frac{HO}{A_bB_a}.$$

Also since $\triangle A_1O_1B_1 \stackrel{+}{\sim} \triangle AHA_b$ (both of them are $\sim \triangle AOB$), we have

$$\frac{A_1O_1}{A_1B_1} = \frac{AH}{AA_b} = \frac{HO}{A_bB_a} = \frac{HO}{A_1B_1} \implies A_1O_1 = HO$$

as desired. \square

Since

$$\sphericalangle (TA_1, TO_1) \stackrel{\text{Claim 5}}{=} \sphericalangle (TO, TA) = 90^\circ + \sphericalangle (TA_1, AA_2) \stackrel{OH \perp AA_2}{=} \sphericalangle (TA_1, OH),$$

we have $O_1T \parallel HO$. Combined with $O_1T = HO$, O_1TOH is a parallelogram. Therefore, using this and *Claim 5*, we have $\sphericalangle H_1O_1T = \sphericalangle TOH = \sphericalangle HO_1T$, which imply that O_1, H_1 and H are collinear as desired.

G8. Let ABC be an equilateral triangle. Points A_1, B_1, C_1 lie inside triangle ABC such that triangle $A_1B_1C_1$ is scalene, $BA_1 = A_1C$, $CB_1 = B_1A$, $AC_1 = C_1B$ and

$$\angle BA_1C + \angle CB_1A + \angle AC_1B = 480^\circ.$$

Lines BC_1 and CB_1 intersect at A_2 ; lines CA_1 and AC_1 intersect at B_2 ; and lines AB_1 and BA_1 intersect at C_2 .

Prove that the circumcircles of triangles AA_1A_2 , BB_1B_2 , CC_1C_2 have two common points.
(U.S.A.)


Solution. Let $\delta_A, \delta_B, \delta_C$ be the circumcircles of $\triangle AA_1A_2$, $\triangle BB_1B_2$, $\triangle CC_1C_2$. The general strategy of the solution is to find two different points having equal power with respect to $\delta_A, \delta_B, \delta_C$.

Claim. A_1 is the circumcentre of A_2BC and cyclic variations.

Proof. Since A_1 lies on the perpendicular bisector of BC and inside $\triangle BA_2C$, it suffices to prove $\angle BA_1C = 2\angle BA_2C$. This follows from

$$\begin{aligned} \angle BA_2C &= \angle A_2BA + \angle BAC + \angle ACA_2 \\ &= \frac{1}{2} ((180^\circ - \angle AC_1B) + (180^\circ - \angle CB_1A)) + 60^\circ \\ &= 240^\circ - \frac{1}{2} (480^\circ - \angle BA_1C) \\ &= \frac{1}{2} \angle BA_1C \end{aligned}$$

□



The circumcentres above give

$$\angle B_1B_2C_1 = \angle B_1B_2A = \angle B_2AB_1 = \angle C_1AC_2 = \angle AC_2C_1 = \angle B_1C_2C_1$$

and so $B_1C_1B_2C_2$ is cyclic. Likewise $C_1A_1C_2A_2$ and $A_1B_1A_2B_2$ are cyclic. Note that hexagon $A_1B_2C_1A_2B_1C_2$ is not cyclic since

$$\angle C_2A_1B_2 + \angle B_2C_1A_2 + \angle A_2B_1C_2 = 480^\circ \neq 360^\circ.$$

Thus we can apply radical axis theorem to the three circles to show that A_1A_2, B_1B_2, C_1, C_2 concur at a point X and this point has equal power with respect to $\delta_A, \delta_B, \delta_C$.

Let the circumcircle of $\triangle A_2BC$ meet δ_A at $A_3 \neq A_2$. Define B_3 and C_3 similarly.

Claim. BCB_3C_3 cyclic.


Proof. Using directed angles

$$\begin{aligned} \angle B C_3 C &= \angle B C_3 C_2 + \angle C_2 C_3 C \\ &= \angle B A C_2 + \angle C_2 C_1 C \\ &= 90^\circ + \angle (C_1 C, A C_2) + \angle C_2 C_1 C && (CC_1 \perp AB) \\ &= 90^\circ + \angle C_1 C_2 B_1. \end{aligned}$$

Similarly $\angle C B_3 B = 90^\circ + \angle B_1 B_2 C_1$. Hence, using $B_1 C_1 B_2 C_2$ cyclic

$$\angle B B_3 C = 90^\circ + \angle C_1 B_2 B_1 = 90^\circ + \angle C_1 C_2 B_1 = \angle B C_3 C$$

as required. □



Similarly CAC_3A_3 and ABA_3B_3 are cyclic. $AC_3BA_3CB_3$ is not cyclic because then AB_2CB_3 cyclic would mean B_2 lies on $\odot ABC$ which is impossible since B_2 lies inside $\triangle ABC$. Thus we can apply radical axis theorem to the three circles to get AA_3, BB_3, CC_3 concur at a point Y which has equal power with respect to $\delta_A, \delta_B, \delta_C$.

We now make some technical observations before finishing.

- Let O be the centre of $\triangle ABC$. We have that

$$\angle BA_1C = 480^\circ - \angle CB_1A - \angle AC_1B > 480^\circ - 180^\circ - 180^\circ = 120^\circ.$$

so A_1 lies inside $\triangle BOC$. We have similar results for B_1, C_1 and thus $\triangle BA_1C, \triangle CB_1A, \triangle AC_1B$ have disjoint interiors. It follows that $A_1B_2C_1A_2B_1C_2$ is a convex hexagon thus X lies on segment A_1A_2 and therefore is inside δ_A .

- Since A_1 is the centre of A_2BC we have that $A_1A_2 = A_1A_3$ so, from cyclic quadrilateral $AA_2A_1A_3$ we get that lines AA_2 and $AA_3 \equiv AY$ are reflections in line AA_1 . As X lies on segment A_1A_2 , the only way $X \equiv Y$ is if A_1 and A_2 both lie on the perpendicular bisector of BC . But this forces B_1 and C_1 to also be reflections in this line meaning $A_1B_1 = A_1C_1$ contradicting the scalene condition.

Summarising, we have distinct points X, Y with equal power with respect to $\delta_A, \delta_B, \delta_C$ thus these circles have a common radical axis. As X lies inside δ_A (and similarly δ_B, δ_C), this radical axis intersects the circles at two points and so $\delta_A, \delta_B, \delta_C$ have two points in common.

Comment. An alternative construction for Y comes by observing that

$$\frac{\sin \angle BAA_2}{\sin \angle A_2AC} = \frac{\frac{A_2B}{AA_2} \sin \angle A_2BA}{\frac{A_2C}{AA_2} \sin \angle ACA_2} = \frac{A_2B}{A_2C} \cdot \frac{\sin \angle C_1BA}{\sin \angle ACB_1} = \frac{\sin \angle B_1CB}{\sin \angle CBC_1} \cdot \frac{\sin \angle C_1BA}{\sin \angle ACB_1}$$

and hence

$$\frac{\sin \angle BAA_2}{\sin \angle A_2AC} \cdot \frac{\sin \angle CBB_2}{\sin \angle B_2BA} \cdot \frac{\sin \angle ACC_2}{\sin \angle C_2CB} = 1$$

so by Ceva's theorem, AA_2, BB_2, CC_2 concur and thus we can construct the isogonal conjugate of this point of concurrency which turns out to be Y .

Number Theory

N1. Determine all positive, composite integers n that satisfy the following property: if the positive divisors of n are $1 = d_1 < d_2 < \dots < d_k = n$, then d_i divides $d_{i+1} + d_{i+2}$ for every $1 \leq i \leq k - 2$.

(Colombia)

Answer: $n = p^r$ is a prime power for some $r \geq 2$.

Solution 1. It is easy to see that such an $n = p^r$ with $r \geq 2$ satisfies the condition as $d_i = p^{i-1}$ with $1 \leq i \leq k = r + 1$ and clearly

$$p^{i-1} \mid p^i + p^{i+1}.$$

Now, let us suppose that there is a positive integer n that satisfies the divisibility condition of the problem and that has two different prime divisors p and q . Without loss of generality, we assume $p < q$ and that they are the two smallest prime divisors of n . Then there is a positive integer j such that

$$d_1 = 1, d_2 = p, \dots, d_j = p^{j-1}, d_{j+1} = p^j, d_{j+2} = q,$$

and it follows that

$$d_{k-j-1} = \frac{n}{q}, d_{k-j} = \frac{n}{p^j}, d_{k-j+1} = \frac{n}{p^{j-1}}, \dots, d_{k-1} = \frac{n}{p}, d_k = n.$$

Thus

$$d_{k-j-1} = \frac{n}{q} \mid d_{k-j} + d_{k-j+1} = \frac{n}{p^j} + \frac{n}{p^{j-1}} = \frac{n}{p^j}(p + 1). \quad (1)$$

This gives $p^j \mid q(p + 1)$, which is a contradiction since $\gcd(p, p + 1) = 1$ and $p \neq q$.

Solution 2. Since $d_i d_{k+1-i} = n$, we have the equivalence:

$$d_{k-i-1} \mid d_{k-i} + d_{k-i+1} \iff \frac{n}{d_{i+2}} \mid \frac{n}{d_{i+1}} + \frac{n}{d_i}.$$

We multiply both sides by $d_i d_{i+1} d_{i+2}$ and cancel the n 's to get

$$d_i d_{i+1} \mid d_i d_{i+2} + d_{i+1} d_{i+2}.$$

Hence,

$$d_i \mid d_{i+1} d_{i+2}. \quad (2)$$

Moreover, by the condition of the problem,

$$d_i \mid d_{i+1}(d_{i+1} + d_{i+2}) = d_{i+1}^2 + d_{i+1} d_{i+2}.$$

Combining this with (2) we get that $d_i \mid d_{i+1}^2$ for all $1 \leq i \leq k - 2$.

Let $d_2 = p$ be the smallest prime divisor of n . By induction on i we prove that $p \mid d_i$ for all $2 \leq i \leq k - 1$. The base case $d_2 = p$ is obvious. Let us suppose that $p \mid d_j$ for some $2 \leq j \leq k - 2$. Then we have that

$$p \mid d_j \mid d_{j+1}^2 \implies p \mid d_{j+1}$$

as p is prime, which completes the induction. This implies that n has to be a prime power, as otherwise there would be another prime q that divides n and we would get that $p \mid q$ which is obviously false.

We finally check that the powers of p satisfy the condition in the statement of the problem as in Solution 1.

Solution 3. We start by proving the following claim:

Claim. $d_i \mid d_{i+1}$ for every $1 \leq i \leq k-1$.

Proof. We prove the Claim by induction on i ; it is trivial for $i = 1$ because $d_1 = 1$. Suppose that $2 \leq i \leq k-1$ and the Claim is true for $i-1$, i.e. $d_{i-1} \mid d_i$. By the induction hypothesis and the problem condition, $d_{i-1} \mid d_i$ and $d_{i-1} \mid d_i + d_{i+1}$, so $d_{i-1} \mid d_{i+1}$.

Now consider the divisors $d_{k-i} = \frac{n}{d_{i+1}}$, $d_{k-i+1} = \frac{n}{d_i}$, $d_{k-i+2} = \frac{n}{d_{i-1}}$. By the problem condition,

$$\frac{d_{k-i+1} + d_{k-i+2}}{d_{k-i}} = \frac{\frac{n}{d_i} + \frac{n}{d_{i-1}}}{\frac{n}{d_{i+1}}} = \frac{d_{i+1}}{d_i} + \frac{d_{i+1}}{d_{i-1}}$$

is an integer. We conclude that $\frac{d_{i+1}}{d_i}$ is an integer, so $d_i \mid d_{i+1}$. \square

By the Claim, n cannot have two different prime divisors because the smallest one would divide the other one. Hence, n must be a power of a prime, and powers of primes satisfy the condition of the problem as we saw in Solution 1.

Solution 4. We present here a more technical way of finishing Solution 1 after obtaining (1). We let $v_p(m)$ denote the p -adic valuation of m . Notice that $v_p(n/q) = v_p(n)$ as $\gcd(p, q) = 1$ and that

$$v_p\left(\frac{n}{p^j}(p+1)\right) = v_p(n) - j$$

as $\gcd(p, p+1) = 1$. But (1) implies

$$v_p(n) = v_p(n/q) \leq v_p\left(\frac{n}{p^j}(p+1)\right) = v_p(n) - j$$

which is a contradiction. Thus n has only one prime divisor as desired.

N2. Determine all pairs (a, p) of positive integers with p prime such that $p^a + a^4$ is a perfect square.

(Bangladesh)

Answer: $(a, p) = (1, 3), (2, 3), (6, 3), (9, 3)$ are all the possible solutions.

Solution. Let $p^a + a^4 = b^2$ for some positive integer b . Then we have

$$p^a = b^2 - a^4 = (b + a^2)(b - a^2).$$

Hence both $b + a^2$ and $b - a^2$ are powers of p .

Let $b - a^2 = p^x$ for some integer x . Then $b + a^2 = p^{a-x}$ and $a - x > x$. Therefore, we have

$$2a^2 = (b + a^2) - (b - a^2) = p^{a-x} - p^x = p^x(p^{a-2x} - 1). \quad (1)$$

We shall consider two cases according to whether $p = 2$ or $p \neq 2$. We let $v_p(m)$ denote the p -adic valuation of m .

Case 1 ($p = 2$): In this case,

$$a^2 = 2^{x-1}(2^{a-2x} - 1) = 2^{2v_2(a)}(2^{a-2x} - 1),$$

where the first equality comes from (1) and the second one from $\gcd(2, 2^{a-2x} - 1) = 1$. So, $2^{a-2x} - 1$ is a square.

If $v_2(a) > 0$, then 2^{a-2x} is also a square. So, $2^{a-2x} - 1 = 0$, and $a = 0$ which is a contradiction.

If $v_2(a) = 0$, then $x = 1$, and $a^2 = 2^{a-2} - 1$. If $a \geq 4$, the right hand side is congruent to 3 modulo 4, thus cannot be a square. It is easy to see that $a = 1, 2, 3$ do not satisfy this condition.

Therefore, we do not get any solutions in this case.

Case 2 ($p \neq 2$): In this case, we have $2v_p(a) = x$. Let $m = v_p(a)$. Then we have $a^2 = p^{2m} \cdot n^2$ for some integer $n \geq 1$. So, $2n^2 = p^{a-2x} - 1 = p^{p^m \cdot n - 4m} - 1$.

We consider two subcases.

Subcase 2-1 ($p \geq 5$): By induction, one can easily prove that $p^m \geq 5^m > 4m$ for all m . Then we have

$$2n^2 + 1 = p^{p^m \cdot n - 4m} > p^{p^m \cdot n - p^m} \geq 5^{5^m \cdot (n-1)} \geq 5^{n-1}.$$

But, by induction, one can easily prove that $5^{n-1} > 2n^2 + 1$ for all $n \geq 3$. Therefore, we conclude that $n = 1$ or 2 . If $n = 1$ or 2 , then $p = 3$, which is a contradiction. So there are no solutions in this subcase.

Subcase 2-2 ($p = 3$): Then we have $2n^2 + 1 = 3^{3^m \cdot n - 4m}$. If $m \geq 2$, one can easily prove by induction that $3^m > 4m$. Then we have

$$2n^2 + 1 = 3^{3^m \cdot n - 4m} > 3^{3^m \cdot n - 3^m} = 3^{3^m \cdot (n-1)} \geq 3^{9(n-1)}.$$

Again, by induction, one can easily prove that $3^{9(n-1)} > 2n^2 + 1$ for all $n \geq 2$. Therefore, we conclude that $n = 1$. Then we have $2 \cdot 1^2 + 1 = 3^{3^m - 4m}$ hence $3 = 3^{3^m - 4m}$. Consequently, we have $3^m - 4m = 1$. The only solution of this equation is $m = 2$ in which case we have $a = 3^m \cdot n = 3^2 \cdot 1 = 9$.

If $m \leq 1$, then there are two possible cases: $m = 0$ or $m = 1$.

- If $m = 1$, then we have $2n^2 + 1 = 3^{3n-4}$. Again, by induction, one can easily prove that $3^{3n-4} > 2n^2 + 1$ for all $n \geq 3$. By checking $n = 1, 2$, we only get $n = 2$ as a solution. This gives $a = 3^m \cdot n = 3^1 \cdot 2 = 6$.
- If $m = 0$, then we have $2n^2 + 1 = 3^n$. By induction, one can easily prove that $3^n > 2n^2 + 1$ for all $n \geq 3$. By checking $n = 1, 2$, we find the solutions $a = 3^0 \cdot 1 = 1$ and $a = 3^0 \cdot 2 = 2$.

Therefore, $(a, p) = (1, 3), (2, 3), (6, 3), (9, 3)$ are all the possible solutions.

N3. For positive integers n and $k \geq 2$ define $E_k(n)$ as the greatest exponent r such that k^r divides $n!$. Prove that there are infinitely many n such that $E_{10}(n) > E_9(n)$ and infinitely many m such that $E_{10}(m) < E_9(m)$.

(Brazil)

Solution 1. We let $v_p(m)$ denote the p -adic valuation of m . By Legendre's Formula we know, for p prime, that $v_p(n!) = \lfloor n/p \rfloor + \lfloor n/p^2 \rfloor + \dots$. We can see that $E_9(n) = \lfloor \frac{v_3(n!)}{2} \rfloor$. Since $v_5(n!) \leq v_2(n!)$ and $E_{10}(n) = \min(v_5(n!), v_2(n!))$, we have $E_{10}(n) = v_5(n!)$.

Let l be a positive integer. Set $n = 5^{2l-1}$. Then we have

$$E_{10}(n) = v_5(n!) = 5^{2l-2} + 5^{2l-3} + \dots + 5 + 1 = \frac{5^{2l-1} - 1}{4} = \frac{n-1}{4}.$$

Since $n = 5^{2l-1} \equiv 2 \pmod{3}$, we have $\lfloor \frac{n}{3} \rfloor = \frac{n-2}{3}$ and it implying

$$v_3(n!) = \left\lfloor \frac{n}{3} \right\rfloor + \left\lfloor \frac{n}{3^2} \right\rfloor + \left\lfloor \frac{n}{3^3} \right\rfloor + \dots < \frac{n-2}{3} + \frac{n}{3^2} + \frac{n}{3^3} + \dots = \frac{n}{2} - \frac{2}{3}.$$

From this we obtain

$$E_9(n) = \left\lfloor \frac{v_3(n!)}{2} \right\rfloor \leq \frac{v_3(n!)}{2} \leq \frac{n}{4} - \frac{1}{3} < \frac{n}{4} - \frac{1}{4} = E_{10}(n).$$

In a similar way, we set now $m = 3^{4l-2}$. Then we have

$$v_3(m!) = 3^{4l-3} + 3^{4l-4} + \dots + 3 + 1 = \frac{3^{4l-2} - 1}{2} = \frac{m-1}{2}.$$

Note that $m = 3^{4l-2} \equiv 1 \pmod{4}$ and hence $E_9(m) = \lfloor \frac{v_3(m!)}{2} \rfloor = \lfloor \frac{m-1}{4} \rfloor = \frac{m-1}{4}$. We also have $m = 3^{4l-2} \equiv 4 \pmod{5}$ implying $\lfloor \frac{m}{5} \rfloor = \frac{m-4}{5}$. Therefore we obtain

$$E_{10}(m) = v_5(m!) = \left\lfloor \frac{m}{5} \right\rfloor + \left\lfloor \frac{m}{5^2} \right\rfloor + \dots < \frac{m-4}{5} + \frac{m}{5^2} + \dots = \frac{m}{4} - \frac{4}{5} < \frac{m}{4} - \frac{1}{4} = E_9(m).$$

We can take infinitely many $n = 5^{2l-1}$ and $m = 3^{4l-2}$ completing the proof.

Solution 2. In the setting of Solution 1, we consider two subsequences:

First, we take $n = 5^{3^{b-1}}$ with $b \geq 2$. Because 5 is not a square modulo 3 and $\varphi(3^b) = 2 \cdot 3^{b-1}$, we have $n \equiv -1 \pmod{3^b}$. Hence,

$$v_3(n!) = \left\lfloor \frac{n}{3} \right\rfloor + \left\lfloor \frac{n}{3^2} \right\rfloor + \dots < \frac{n-2}{3} + \frac{n-8}{9} + \dots + \frac{n-(3^b-1)}{3^b} + \frac{n}{3^{b+1}} + \dots < \frac{n}{2} - b + \frac{1}{2},$$

and $E_{10}(n) = \frac{n-1}{4} > \frac{n+1-2b}{4} > E_9(n)$.

In the same way, for $m = 3^{2 \cdot 5^{b-1}} \equiv -1 \pmod{5^b}$ with $b \geq 2$,

$$E_{10}(m) = \left\lfloor \frac{m}{5} \right\rfloor + \left\lfloor \frac{m}{5^2} \right\rfloor + \dots < \frac{m-4}{5} + \frac{m-24}{25} + \dots + \frac{m-(5^b-1)}{5^b} + \frac{m}{5^{b+1}} + \dots < \frac{m}{4} - b + \frac{1}{4},$$

and $E_9(m) = \frac{m-1}{4} > E_{10}(m)$ holds.

Comment. From Solution 2 we can see that for any positive real B , there exist infinitely many positive integers m and n such that $E_{10}(n) - E_9(n) > B$ and $E_{10}(m) - E_9(m) < -B$.

N4.

Let $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n$ be $2n$ positive integers such that the $n + 1$ products

$$\begin{aligned} & a_1 a_2 a_3 \cdots a_n, \\ & b_1 a_2 a_3 \cdots a_n, \\ & b_1 b_2 a_3 \cdots a_n, \\ & \vdots \\ & b_1 b_2 b_3 \cdots b_n \end{aligned}$$

form a strictly increasing arithmetic progression in that order. Determine the smallest positive integer that could be the common difference of such an arithmetic progression.

(Canada)

Answer: The smallest common difference is $n!$.

Solution 1. The condition in the problem is equivalent to

$$D = (b_1 - a_1)a_2a_3 \cdots a_n = b_1(b_2 - a_2)a_3a_4 \cdots a_n = \cdots = b_1b_2 \cdots b_{n-1}(b_n - a_n),$$

where D is the common difference. Since the progression is strictly increasing, $D > 0$, hence $b_i > a_i$ for every $1 \leq i \leq n$. Individually, these equalities simplify to

$$(b_i - a_i)a_{i+1} = b_i(b_{i+1} - a_{i+1}) \text{ for every } 1 \leq i \leq n - 1. \quad (1)$$

If $g_i := \gcd(a_i, b_i) > 1$ for some $1 \leq i \leq n$, then we can replace a_i with $\frac{a_i}{g_i}$ and b_i with $\frac{b_i}{g_i}$ to get a smaller common difference. Hence we may assume $\gcd(a_i, b_i) = 1$ for every $1 \leq i \leq n$.

Then, we have $\gcd(b_i - a_i, b_i) = \gcd(a_i, b_i) = 1$ and $\gcd(a_{i+1}, b_{i+1} - a_{i+1}) = \gcd(a_{i+1}, b_{i+1}) = 1$ for every $1 \leq i \leq n - 1$. The equality (1) implies $a_{i+1} = b_i$ and $b_i - a_i = b_{i+1} - a_{i+1}$. Thus,

$$a_1, \quad b_1 = a_2, \quad b_2 = a_3, \quad \dots, \quad b_{n-1} = a_n, \quad b_n$$

is an arithmetic progression with positive common difference. Since $a_1 \geq 1$, we have $a_i \geq i$ for every $1 \leq i \leq n$, so

$$D = (b_1 - a_1)a_2a_3 \cdots a_n \geq 1 \cdot 2 \cdot 3 \cdots n = n!$$

Equality is achieved when $b_i - a_i = 1$ for $1 \leq i \leq n$ and $a_1 = 1$, i.e. $a_i = i$ and $b_i = i + 1$ for every $1 \leq i \leq n$. Indeed, it is straightforward to check that these integers produce an arithmetic progression with common difference $n!$.

Solution 2 (Variant of Solution 1). Similarly to Solution 1, we may assume $\gcd(a_i, b_i) = 1$ for every $1 \leq i \leq n$.

Denote by p_1, p_2, \dots, p_{n+1} the sequence obtained as the product in the problem statement. Then we have $\frac{p_{i+1}}{p_i} = \frac{b_i}{a_i} > 1$, so $b_i > a_i$. Since p_1, p_2, \dots, p_{n+1} is an arithmetic progression, we have $p_{i+2} = 2p_{i+1} - p_i$ hence

$$2 - \frac{a_i}{b_i} = \frac{2b_i - a_i}{b_i} = \frac{2p_{i+1} - p_i}{p_{i+1}} = \frac{p_{i+2}}{p_{i+1}} = \frac{b_{i+1}}{a_{i+1}}.$$

But since the fractions on the left-hand side and the right-hand side are both irreducible, we conclude that $b_i = a_{i+1}$, so $2 - \frac{a_i}{a_{i+1}} = \frac{a_{i+2}}{a_{i+1}}$. Then we have $a_i + a_{i+2} = 2a_{i+1}$, which means that a_1, a_2, \dots, a_n is an arithmetic progression with positive common difference.

We conclude as in Solution 1.

Solution 3. (The following solution is purely algebraic: it does not involve considerations on greatest common divisors.)

We retake Solution 1 from (1). Then we have

$$\frac{a_{i+1}}{b_{i+1} - a_{i+1}} = \frac{b_i}{b_i - a_i} = 1 + \frac{a_i}{b_i - a_i}.$$

So, for $1 \leq i \leq n$,

$$\frac{a_i}{b_i - a_i} = \frac{a_1}{b_1 - a_1} + (i - 1).$$

Then

$$a_i \geq \frac{a_i}{b_i - a_i} = \frac{a_1}{b_1 - a_1} + (i - 1) > i - 1.$$

since $b_i - a_i \geq 1$ and $b_1 - a_1 > 0$. As a_i is an integer, we have $a_i \geq i$.

We again conclude as in Solution 1.

N5. Let $a_1 < a_2 < a_3 < \cdots$ be positive integers such that a_{k+1} divides $2(a_1 + a_2 + \cdots + a_k)$ for every $k \geq 1$. Suppose that for infinitely many primes p , there exists k such that p divides a_k . Prove that for every positive integer n , there exists k such that n divides a_k .

(Netherlands)

Solution. For every $k \geq 2$ define the quotient $b_k = 2(a_1 + \cdots + a_{k-1})/a_k$, which must be a positive integer. We first prove the following properties of the sequence (b_k) :

Claim 1. We have $b_{k+1} \leq b_k + 1$ for all $k \geq 2$.

Proof. By subtracting $b_k a_k = 2(a_1 + \cdots + a_{k-1})$ from $b_{k+1} a_{k+1} = 2(a_1 + \cdots + a_k)$, we find that $b_{k+1} a_{k+1} = b_k a_k + 2a_k = (b_k + 2)a_k$. From $a_k < a_{k+1}$ it follows that $b_k + 2 > b_{k+1}$. \square

Claim 2. The sequence (b_k) is unbounded.

Proof. We start by rewriting $b_{k+1} a_{k+1} = (b_k + 2)a_k$ as

$$a_{k+1} = a_k \cdot \frac{b_k + 2}{b_{k+1}} \implies a_{k+1} \mid a_k (b_k + 2).$$

If the sequence (b_k) were bounded, say by some positive integer B , then the prime factors of the terms of the sequence (a_k) could only be primes less than or equal to $B + 2$ or those dividing a_1 or a_2 , which contradicts the property in the statement of the problem. \square

Consider now an arbitrary positive integer n . We assume $n > b_2$, otherwise we replace n by an arbitrary multiple of n that is bigger than b_2 . By Claim 2, there exists k such that $b_{k+1} \geq n$. Consider the smallest such k . From Claim 1, it follows that we must have $b_k = n - 1$ and $b_{k+1} = n$ (we assumed $n > b_2$ to ensure that $k \geq 2$). We now find that

$$a_{k+1} = a_k \cdot \frac{b_k + 2}{b_{k+1}} = a_k \cdot \frac{n + 1}{n}.$$

Because n and $n + 1$ are coprime, this immediately implies that a_k is divisible by n .

Comment. For c a positive integer, the sequence $a_k = ck$ satisfies the conditions of the problem. Another example is

$$a_1 = 1, \quad a_2 = 2, \quad a_k = 3(k - 1) \text{ for } k \geq 3.$$

N6.

A sequence of integers a_0, a_1, a_2, \dots is called *kawaii*, if $a_0 = 0, a_1 = 1$, and, for any positive integer n , we have

$$(a_{n+1} - 3a_n + 2a_{n-1})(a_{n+1} - 4a_n + 3a_{n-1}) = 0.$$

An integer is called *kawaii* if it belongs to a *kawaii* sequence.

Suppose that two consecutive positive integers m and $m+1$ are both *kawaii* (not necessarily belonging to the same *kawaii* sequence). Prove that 3 divides m , and that $m/3$ is *kawaii*.

(China)

Solution 1. We start by rewriting the condition in the problem as:

$$a_{n+1} = 3a_n - 2a_{n-1}, \text{ or } a_{n+1} = 4a_n - 3a_{n-1}.$$

We have $a_{n+1} \equiv a_n$ or $a_{n-1} \pmod{2}$ and $a_{n+1} \equiv a_{n-1}$ or $a_n \pmod{3}$ for all $n \geq 1$. Now, since $a_0 = 0$ and $a_1 = 1$, we have that $a_n \equiv 0, 1 \pmod{3}$ for all $n \geq 0$. Since m and $m+1$ are *kawaii* integers, then necessarily $m \equiv 0 \pmod{3}$.

We also observe that $a_2 = 3$ or $a_2 = 4$. Moreover,

(1) If $a_2 = 3$, then $a_n \equiv 1 \pmod{2}$ for all $n \geq 1$ since $a_1 \equiv a_2 \equiv 1 \pmod{2}$.

(2) If $a_2 = 4$, then $a_n \equiv 1 \pmod{3}$ for all $n \geq 1$ since $a_1 \equiv a_2 \equiv 1 \pmod{3}$.

Since $m \equiv 0 \pmod{3}$, any *kawaii* sequence containing m does not satisfy (2), so it must satisfy (1). Hence, m is odd and $m+1$ is even.

Take a *kawaii* sequence (a_n) containing $m+1$. Let $t \geq 2$ be such that $a_t = m+1$. As (a_n) does not satisfy (1), it must satisfy (2). Then $a_n \equiv 1 \pmod{3}$ for all $n \geq 1$. We define the sequence $a'_n = (a_{n+1} - 1)/3$. This is a *kawaii* sequence: $a'_0 = 0, a'_1 = 1$ and for all $n \geq 1$,

$$(a'_{n+1} - 3a'_n + 2a'_{n-1})(a'_{n+1} - 4a'_n + 3a'_{n-1}) = (a_{n+2} - 3a_{n+1} + 2a_n)(a_{n+2} - 4a_{n+1} + 3a_n)/9 = 0.$$

Finally, we notice that the term $a'_{t-1} = m/3$ which implies that $m/3$ is *kawaii*.

Solution 2. We start by proving the following:

Claim 1. We have $a_n \equiv 0, 1 \pmod{3}$ for all $n \geq 0$.

Proof. We have $a_{n+1} = 3a_n - 2a_{n-1} = 3(a_n - a_{n-1}) + a_{n-1}$ or $a_{n+1} = 4a_n - 3a_{n-1} = 3(a_n - a_{n-1}) + a_n$, so $a_{n+1} \equiv a_n$ or $a_{n-1} \pmod{3}$, and since $a_0 = 0$ and $a_1 = 1$ the result follows. \square

Hence if m and $m+1$ are *kawaii*, then necessarily $m \equiv 0 \pmod{3}$.

Claim 2. An integer ≥ 2 is *kawaii* if and only if it can be written as $1 + b_2 + \dots + b_n$ for some $n \geq 2$ with $b_i = 2^{r_i}3^{s_i}$ satisfying $r_i + s_i = i - 1$ for $i = 2, \dots, n$ and $b_i \mid b_{i+1}$ for all $i = 2, \dots, n - 1$.

Proof. For a *kawaii* sequence (a_n) , we can write $a_{n+1} = 3a_n - 2a_{n-1} = a_n + 2(a_n - a_{n-1})$ or $a_{n+1} = 4a_n - 3a_{n-1} = a_n + 3(a_n - a_{n-1})$, so $a_{n+1} - a_n = 2(a_n - a_{n-1})$ or $3(a_n - a_{n-1})$. Hence, $a_n = 1 + b_2 + \dots + b_n$ where $b_2 = 2$ or 3 and $b_{i+1} = 2b_i$ or $3b_i$.

Conversely, given a number that can be written in that way, we consider any sequence given by $a_0 = 0, a_1 = 1$ and $a_i = 1 + b_2 + \dots + b_i$ for $2 \leq i \leq n$ and a_i given by the *kawaii* condition for $i \geq n+1$. This defines a *kawaii* sequence containing the given number as a_n . \square

Let us suppose that m and $m+1$ are *kawaii*, then they belong to some *kawaii* sequences and we can write them as in Claim 2 as $m = 1 + 2 + \dots + 2^\ell + 2^\ell \cdot 3 \cdot A$ and $m+1 = 1 + 2 + \dots + 2^{\ell'} + 2^{\ell'} \cdot 3 \cdot A'$ where ℓ is odd and ℓ' is even because of modulo 3 reasons. Since $m+1 \equiv m \pmod{2^{\min(\ell, \ell')}}$, we have $\min(\ell, \ell') = 0$, so $\ell' = 0$.

Then $m+1 = 1 + b_2 + \dots + b_j$ for some b_i 's as in Claim 2 with $b_2 = 3$ and $b_i \mid b_{i+1}$: so with $3 \mid b_i$ for all $i = 2, \dots, j$. Then $\frac{m}{3} = 1 + b'_1 + \dots + b'_{j-1}$ with $b'_i = \frac{b_{i+1}}{3}$ as in Claim 2 and $\frac{m}{3}$ is a *kawaii* integer.

Solution 3. (This solution is just a different combination of the ideas in Solutions 1 and 2) We first prove that, in a kawaii sequence a_0, a_1, a_2, \dots , every term a_t with $t \geq 0$ is congruent to 0 or 1 modulo 3.

For $n \geq 1$, put $b_n = a_n - a_{n-1}$. We have

$$a_t = a_0 + \sum_{k=1}^t (a_k - a_{k-1}) = \sum_{k=1}^t b_k. \quad (*)$$

Note that

$$a_{n+1} - 3a_n + 2a_{n-1} = b_{n+1} - 2b_n \quad \text{and} \quad a_{n+1} - 4a_n + 3a_{n-1} = b_{n+1} - 3b_n.$$

The conditions on the b_i 's for defining a kawaii sequence are

$$b_1 = a_1 - a_0 = 1, \quad \text{and} \quad \frac{b_{n+1}}{b_n} \in \{2, 3\} \quad \text{for} \quad n \geq 1.$$

1. If we have $\frac{b_{n+1}}{b_n} = 2$ for any n with $1 \leq n \leq t-1$, then $(*)$ implies that

$$a_t = \sum_{k=1}^t 2^{k-1} = 2^t - 1 \equiv 0, 1 \pmod{3}.$$

2. If there exists some integer s with $2 \leq s \leq t-1$ such that

$$\frac{b_2}{b_1} = \frac{b_3}{b_2} = \dots = \frac{b_s}{b_{s-1}} = 2, \quad \frac{b_{s+1}}{b_s} = 3,$$

it implies that $3 \mid b_n$ for any $n \geq s+1$. Similarly to the argument in (1), we obtain

$$a_t \equiv \sum_{k=1}^s b_k \equiv 0, 1 \pmod{3}.$$

3. If $\frac{b_2}{b_1} = b_2 = 3$, we have $3 \mid b_n$ for any $n \geq 2$, and hence $a_t \equiv 1 \pmod{3}$.

Combining these, we have proved that $a_t \equiv 0, 1 \pmod{3}$.

We next prove that no positive kawaii integer is divisible by both 2 and 3. If $b_2 = 2$ for some kawaii sequence, then $2 \mid b_n$ and $a_n \equiv 1 \pmod{2}$ for all $n \geq 2$ in it. If $b_2 = 3$ in some kawaii sequence, then $3 \mid b_n$ and $a_n \equiv 1 \pmod{3}$ for all $n \geq 2$ in it.

Now, consider the original problem. Since m and $m+1$ are both kawaii integer, it means

$$m \equiv 0, 1 \pmod{3}, \quad \text{and} \quad m+1 \equiv 0, 1 \pmod{3},$$

and hence we easily obtain $3 \mid m$. Since a kawaii integer m is divisible by 3, m must be odd, and hence $m+1$ is even. Take a kawaii sequence a_0, a_1, a_2, \dots containing $m+1$ as a_t . The fact that $m+1$ is even implies that $b_2 = 3$ and so $3 \mid b_n$ for all $n \geq 2$ in this sequence. Set $b'_n = \frac{b_{n+1}}{3}$ for $n \geq 1$. Thus $b'_1 = \frac{b_2}{3} = 1$, and $\frac{b'_{n+1}}{b'_n} = \frac{b_{n+2}}{b_{n+1}} \in \{2, 3\}$ for all $n \geq 1$. Define $a'_0 = 0$

and $a'_n = \sum_{k=1}^n b'_k$ for $n \geq 1$, then a'_0, a'_1, a'_2, \dots is a kawaii sequence. Now,

$$a'_{t-1} = \sum_{k=1}^{t-1} b'_k = \frac{1}{3} \sum_{k=2}^t b_k = \frac{1}{3} \left(-b_1 + \sum_{k=1}^t b_k \right) = \frac{1}{3} (-1 + a_t) = \frac{m}{3}.$$

This means that $\frac{m}{3}$ is a kawaii integer.

Comment. There are infinitely many positive integers m such that $m, m+1, m/3$ are kawaii. To see this, let $k \geq 1$ be a kawaii integer. Then $2k+1$ and $3k+1$ are kawaii by Claim 2 in Solution 2, and $3(2k+1)+1 = 6k+4$ and $2(3k+1)+1 = 6k+3$ are also kawaii.

N7. Let a, b, c, d be positive integers satisfying

$$\frac{ab}{a+b} + \frac{cd}{c+d} = \frac{(a+b)(c+d)}{a+b+c+d}.$$

Determine all possible values of $a+b+c+d$.

(Netherlands)

Answer: The possible values are the positive integers that are not square-free.

Solution.

First, note that if we take $a = \ell$, $b = k\ell$, $c = k\ell$, $d = k^2\ell$ for some positive integers k and ℓ , then we have

$$\frac{ab}{a+b} + \frac{cd}{c+d} = \frac{k\ell^2}{\ell+k\ell} + \frac{k^3\ell^2}{k\ell+k^2\ell} = \frac{k\ell}{k+1} + \frac{k^2\ell}{k+1} = k\ell$$

and

$$\frac{(a+b)(c+d)}{a+b+c+d} = \frac{(\ell+k\ell)(k\ell+k^2\ell)}{\ell+k\ell+k\ell+k^2\ell} = \frac{k(k+1)^2\ell^2}{\ell(k+1)^2} = k\ell,$$

so that

$$\frac{ab}{a+b} + \frac{cd}{c+d} = k\ell = \frac{(a+b)(c+d)}{a+b+c+d}.$$

This means that $a+b+c+d = \ell(1+2k+k^2) = \ell(k+1)^2$ can be attained. We conclude that all non-square-free positive integers can be attained.

Now, we will show that if

$$\frac{ab}{a+b} + \frac{cd}{c+d} = \frac{(a+b)(c+d)}{a+b+c+d}$$

then $a+b+c+d$ is not square-free. We argue by contradiction. Suppose that $a+b+c+d$ is square-free, and note that after multiplying by $(a+b)(c+d)(a+b+c+d)$, we obtain

$$(ab(c+d) + cd(a+b))(a+b+c+d) = (a+b)^2(c+d)^2. \quad (1)$$

A prime factor of $a+b+c+d$ must divide $a+b$ or $c+d$, and therefore divides both $a+b$ and $c+d$. Because $a+b+c+d$ is square-free, the fact that every prime factor of $a+b+c+d$ divides $a+b$ implies that $a+b+c+d$ itself divides $a+b$. Because $a+b < a+b+c+d$, this is impossible. So $a+b+c+d$ cannot be square-free.

Comment 1. Another way to conclude after obtaining (1) is by observing that

$$(a+b)^2(c+d)^2 \equiv (a+b)^4 \pmod{a+b+c+d}$$

Hence $a+b+c+d \mid (a+b)^4$. But if $a+b+c+d$ is square-free, this forces $a+b+c+d \mid a+b$, which is clearly a contradiction.

Comment 2. It seems difficult to characterise all quadruples (a, b, c, d) that satisfy the equality in the problem. Many of them, including those used in the solution, are of the general form $(a, b, c, d) = (xy^2, xyz, xyz, xz^2)$ for some positive integers x, y , and z , but there are more solutions than that, such as $(a, b, c, d) = (2, 7, 8, 10)$ or $(a, b, c, d) = (13, 14, 16, 38)$.

N8. Let $\mathbb{Z}_{>0}$ be the set of positive integers. Determine all functions $f: \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ such that

$$f^{bf(a)}(a+1) = (a+1)f(b)$$

holds for all $a, b \in \mathbb{Z}_{>0}$, where $f^k(n) = f(f(\cdots f(n)\cdots))$ denotes the composition of f with itself k times.

(Taiwan)

Answer: The only function satisfying the condition is $f(n) = n + 1$ for all $n \in \mathbb{Z}_{>0}$.

Let $P(a, b)$ be the equality in the statement.

Solution 1. We divide the solution into 5 steps.

Step 1. (f is injective)

Claim 1. For any $a \geq 2$, the set $\{f^n(a) \mid n \in \mathbb{Z}_{>0}\}$ is infinite.

Proof. First, we have $f^{f(a)}(a+1) \stackrel{P(a,1)}{=} (a+1)f(1)$. Varying a , we see that $f(\mathbb{Z}_{>0})$ is infinite. Next, we have $f^{bf(a-1)}(a) \stackrel{P(a-1,b)}{=} af(b)$. So, varying b , $f^{bf(a-1)}(a)$ takes infinitely many values. \square

Claim 2. For any $a \geq 2$ and $n \in \mathbb{Z}_{>0}$, we have $f^n(a) \neq a$.

Proof. Otherwise we would get a contradiction with Claim 1. \square

Assume $f(b) = f(c)$ for some $b < c$. Then we have

$$\begin{aligned} (a+1)f(c) &\stackrel{P(a,c)}{=} f^{cf(a)}(a+1) \\ &= f^{(c-b)f(a)}(f^{bf(a)}(a+1)) \\ &\stackrel{P(a,b)}{=} f^{(c-b)f(a)}((a+1)f(b)) \\ &= f^{(c-b)f(a)}((a+1)f(c)), \end{aligned}$$

which contradicts Claim 2. So, f is injective.

Step 2. ($f(\mathbb{Z}_{>0}) = \mathbb{Z}_{\geq 2}$)

Claim 3. 1 is not in the range of f .

Proof. If $f(b) = 1$, then $f^{f(a)}(a+1) = a+1$ by $P(a,1)$, which contradicts Claim 2. \square

We say that a is a *descendant* of b if $f^n(b) = a$ for some $n \in \mathbb{Z}_{>0}$.

Claim 4. For any $a, b \geq 1$, both of the following cannot happen at the same time:

- a is a descendant of b ;
- b is a descendant of a .

Proof. If both of the above hold, then $a = f^m(b)$ and $b = f^n(a)$ for some $m, n \in \mathbb{Z}_{>0}$. Then $a = f^{m+n}(a)$, which contradicts Claim 2. \square

Claim 5. For any $a, b \geq 2$, exactly one of the following holds:

- a is a descendant of b ;
- b is a descendant of a ;
- $a = b$.

Proof. For any $c \geq 2$, taking $m = f^{cf(a-1)-1}(a)$ and $n = f^{cf(b-1)-1}(b)$, we have

$$f(m) = f^{cf(a-1)}(a) \stackrel{P(a-1,c)}{=} af(c) \text{ and } f(n) = f^{cf(b-1)}(b) \stackrel{P(b-1,c)}{=} bf(c).$$

Hence

$$f^{nf(a-1)}(a) \stackrel{P(a-1,n)}{=} af(n) = abf(c) = bf(m) \stackrel{P(a-1,m)}{=} f^{mf(b-1)}(b).$$

The assertion then follows from the injectivity of f and Claim 2. \square

Now, we show that any $a \geq 2$ is in the range of f . Let $b = f(1)$. If $a = b$, then a is in the range of f . If $a \neq b$, either a is a descendant of b , or b is a descendant of a by Claim 5. If b is a descendant of a , then $b = f^n(a)$ for some $n \in \mathbb{Z}_{>0}$, so $1 = f^{n-1}(a)$. Then, by Claim 3, we have $n = 1$, so $1 = a$, which is absurd. So, a is a descendant of b . In particular, a is in the range of f . Thus, $f(\mathbb{Z}_{>0}) = \mathbb{Z}_{\geq 2}$.

Step 3. ($f(1) = 2$)

Claim 6. Let $a, n \geq 2$, then na is a descendant of a .

Proof. We write $n = f(m)$ by Step 2. We have $na = f(m)a \stackrel{P(a-1,m)}{=} f^{mf(a-1)}(a)$, which shows na is a descendant of a . \square

By Claim 6, all even integers ≥ 4 are descendants of 2. Hence $2 = f(2k+1)$ for some $k \geq 0$.

Next, we show $f(2k+1) \geq f(1)$, which implies $f(1) = 2$. It trivially holds if $k = 0$. If $k \geq 1$, let n be the integer such that $f^n(2) = 2k+2$. For any $b > n/f(1)$, we have

$$f^{bf(1)-n}(2k+2) = f^{bf(1)}(2) \stackrel{P(1,b)}{=} 2f(b) \text{ and } f^{bf(2k+1)}(2k+2) \stackrel{P(2k+1,b)}{=} (2k+2)f(b).$$

By Claim 6, $(2k+2)f(b)$ is a descendant of $2f(b)$. By Claim 2, we have $bf(2k+1) > bf(1) - n$. By taking b large enough, we conclude $f(2k+1) \geq f(1)$.

Step 4. ($f(2) = 3$ and $f(3) = 4$) From $f(1) = 2$ and $P(1, b)$, we have $f^{2b}(2) = 2f(b)$. So taking $b = 1$, we obtain $f^2(2) = 2f(1) = 4$; and taking $b = f(2)$, we have $f^{2f(2)}(2) = 2f^2(2) = 8$. Hence, $f^{2f(2)-2}(4) = f^{2f(2)}(2) = 8$ and $f^{f(3)}(4) \stackrel{P(3,1)}{=} 8$ give $f(3) = 2f(2) - 2$.

Claim 7. For any $m, n \in \mathbb{Z}_{>0}$, if $f(m)$ divides $f(n)$, then $m \leq n$.

Proof. If $f(m) = f(n)$, the assertion follows from the injectivity of f . If $f(m) < f(n)$, by $P(a, m)$, $P(a, n)$ and Claim 6, we have that $f^{nf(a)}(a+1)$ is a descendant of $f^{mf(a)}(a+1)$ for any $a \in \mathbb{Z}_{>0}$. So $mf(a) < nf(a)$, and $m < n$. \square

By Claim 7, every possible divisor of $f(2)$ is in $\{1, f(1) = 2, f(2)\}$. Thus $f(2)$ is an odd prime or $f(2) = 4$. Since $f^2(2) = 4$, we have $f(2) \neq 4$, and hence $f(2)$ is an odd prime. We set $p = f(2)$.

Now, $f(3) = 2f(2) - 2 = 2(p-1)$. Since $p-1$ divides $f(3)$, we have $p-1 \in \{1, f(1) = 2, f(2) = p\}$ by Claim 7, so $p-1 = 2$. Thus, $f(2) = p = 3$ and $f(3) = 2(p-1) = 4$.

Step 5. ($f(n) = n+1$)

Claim 8. For any $b \geq 1$, $f(2f(b)-1) = 2b+2$.

Proof. Since $f^2(2) = 4$, we have $f^{2b-2}(4) = f^{2b}(2) = 2f(b)$, so

$$f^{f(2f(b)-1)+2b-2}(4) = f^{f(2f(b)-1)}(2f(b)) \stackrel{P(2f(b)-1,1)}{=} 4f(b) \stackrel{P(3,b)}{=} f^{4b}(4),$$

which gives us $f(2f(b)-1) = 2b+2$. \square

Finally, we prove $f(n) = n+1$ by induction on n . Suppose $f(n) = n+1$ for all $1 \leq n \leq 2b+1$. Replace b by $b+1$ in $f(2f(b)-1) = 2b+2$ to get

$$f(2b+3) = f(2f(b+1)-1) = 2(b+1)+2 = 2b+4.$$

By induction hypothesis, we have $f^b(b+2) = 2b+2$. Hence

$$f(f(2b+2)) = f^{b+2}(b+2) = f^{f(b+1)}(b+2) \stackrel{P(b+1,1)}{=} 2(b+2) = f(2b+3).$$

By injectivity, $f(2b+2) = 2b+3$. Then $f(n) = n+1$ for all $n \in \mathbb{Z}_{>0}$, which is indeed a solution.

Solution 2. In the same way as Steps 1-2 of Solution 1, we have that f is injective and $f(\mathbb{Z}_{>0}) = \mathbb{Z}_{\geq 2}$.

We first note that Claim 2 in Solution 1 is also true for $a = 1$.

Claim 2'. For any $a, n \in \mathbb{Z}_{>0}$, we have $f^n(a) \neq a$.

Proof. If $a \geq 2$, the assertion was proved in Claim 2 in Solution 1. If $a = 1$, we have that 1 is not in the range of f by Claim 3 in Solution 1. So, $f^n(1) \neq 1$ for every $n \in \mathbb{Z}_{>0}$. \square

For any $a, b \in \mathbb{Z}_{>0}$, we have

$$f^{bf(f(a)-1)+1}(a) = f^{bf(f(a)-1)}(f(a)) \stackrel{P(f(a)-1,b)}{=} f(a)f(b).$$

Since the right-hand side is symmetric in a, b , we have

$$f^{bf(f(a)-1)+1}(a) = f(a)f(b) = f^{af(f(b)-1)+1}(b).$$

Since f is injective, we have $f^{bf(f(a)-1)}(a) = f^{af(f(b)-1)}(b)$. We set $g(n) = f(f(n)-1)$. Then we have $f^{bg(a)}(a) = f^{ag(b)}(b)$ for any $a, b \in \mathbb{Z}_{>0}$. We set $n_{a,b} = bg(a) - ag(b)$. Then, for sufficiently large n , we have $f^{n+n_{a,b}}(a) = f^n(b)$. For any $a, b, c \in \mathbb{Z}_{>0}$ and sufficiently large n , we have

$$f^{n+n_{a,b}+n_{b,c}+n_{c,a}}(a) = f^n(a).$$

By Claim 2' above, we have $n_{a,b} + n_{b,c} + n_{c,a} = 0$, so

$$(a-b)g(c) + (b-c)g(a) + (c-a)g(b) = 0.$$

Taking $(a, b, c) = (n, n+1, n+2)$, we have $g(n+1) - g(n) = g(n+2) - g(n+1)$. So, $\{g(n)\}_{n \geq 1}$ is an arithmetic progression.

There exist $C, D \in \mathbb{Z}$ such that $g(n) = f(f(n)-1) = Cn + D$ for all $n \in \mathbb{Z}_{>0}$. By Step 2 of Solution 1, we have $f(\mathbb{Z}_{>0}) = \mathbb{Z}_{\geq 2}$, so $C = 1$. Since $2 = \min_{n \in \mathbb{Z}_{>0}} \{f(f(n)-1)\}$, we have $D = 1$.

Thus, $g(n) = f(f(n)-1) = n+1$ for all $n \geq 1$. For any $a, b \in \mathbb{Z}_{>0}$, we have $f^{b(a+1)}(a) = f^{a(b+1)}(b)$. By the injectivity of f , we have $f^b(a) = f^a(b)$. For any $n \in \mathbb{Z}_{>0}$, taking $(a, b) = (1, n)$, we have $f^n(1) = f(n)$, so $f^{n-1}(1) = n$ again by the injectivity of f . For any $n \geq 1$, we have $f(n) = f(f^{n-1}(1)) = f^n(1) = n+1$.

Solution 3. The following is another way of finishing Solution 2 after Claim 2' and having introduced $g(n) = f(f(n)-1)$. For $a, b \in \mathbb{Z}_{>0}$ satisfying $b = f^k(a)$, we have

$$\begin{aligned} f^{bg(a)}(f(a)) &= f^{bf(f(a)-1)}(f(a)) \stackrel{P(f(a)-1,b)}{=} f(a)f(b) \stackrel{P(f(b)-1,a)}{=} f^{af(f(b)-1)}(f(b)) \\ &= f^{ag(b)+k}(f(a)). \end{aligned}$$

By Claim 2' in Solution 2, we have $bg(a) = ag(b) + k$, so $f^k(a) \cdot g(a) = a \cdot g(f^k(a)) + k$. Therefore, for any $n \geq 0$, we put $a_n = f^n(1)$. We have

$$\begin{cases} a_{n+1} \cdot g(a_n) = a_n \cdot g(a_{n+1}) + 1 \\ a_{n+2} \cdot g(a_{n+1}) = a_{n+1} \cdot g(a_{n+2}) + 1 \\ a_{n+2} \cdot g(a_n) = a_n \cdot g(a_{n+2}) + 2. \end{cases}$$

Then we have

$$\begin{aligned} a_n \cdot a_{n+1} \cdot g(a_{n+2}) + 2a_{n+1} &= a_{n+1} \cdot a_{n+2} \cdot g(a_n) \\ &= a_{n+2} (a_n \cdot g(a_{n+1}) + 1) \\ &= a_n \cdot a_{n+2} \cdot g(a_{n+1}) + a_{n+2} \\ &= a_n (a_{n+1} \cdot g(a_{n+2}) + 1) + a_{n+2} \\ &= a_n \cdot a_{n+1} \cdot g(a_{n+2}) + a_n + a_{n+2}. \end{aligned}$$

From these, we have $2a_{n+1} = a_n + a_{n+2}$. Thus (a_n) is an arithmetic progression, and we have $a_n = f^n(1) = Cn + D$ for some $C, D \in \mathbb{Z}$.

By Step 2 in Solution 1, any $a \in \mathbb{Z}_{>0}$ is a descendant of 1, and $f(\mathbb{Z}_{>0}) = \mathbb{Z}_{\geq 2}$. Hence $D = 1$ and $C = 1$, and so $f^n(1) = n + 1$. For any $n \geq 1$, we have $f^{n-1}(1) = n$, so $f(n) = f(f^{n-1}(1)) = f^n(1) = n + 1$.

Solution 4.

We provide yet another (more technical) solution assuming Step 1 and Step 2 of Solution 1.

By Claim 5 in Solution 1, every $a \geq 2$ is a descendant of 1. Let g and h be the functions on $\mathbb{Z}_{\geq 2}$ such that $f^{g(a)}(1) = a$ and $h(a) = f(a - 1)$. Then, $g: \mathbb{Z}_{\geq 2} \rightarrow \mathbb{Z}_{\geq 1}$ and $h: \mathbb{Z}_{\geq 2} \rightarrow \mathbb{Z}_{\geq 2}$ are bijections. The equation $P(a, b)$ can be rewritten as

$$g(ah(b)) = g(a) + (b - 1)h(a).$$

Consider the set $S_a = g(a \cdot \mathbb{Z}_{>0})$. Since h is a bijection onto $\mathbb{Z}_{\geq 2}$, we have

$$S_a = g(a) + h(a) \cdot \mathbb{Z}_{\geq 0}.$$

Consider the intersection $S_a \cap S_b = S_{\text{lcm}(a, b)}$. If we put $c = \text{lcm}(a, b)$, this gives

$$(g(a) + h(a) \cdot \mathbb{Z}_{\geq 0}) \cap (g(b) + h(b) \cdot \mathbb{Z}_{\geq 0}) = g(c) + h(c) \cdot \mathbb{Z}_{\geq 0}.$$

Then we have $h(c) = \text{lcm}(h(a), h(b))$ since the left hand side must be of the form $m + \text{lcm}(h(a), h(b)) \cdot \mathbb{Z}_{\geq 0}$ for some m .

If b is a multiple of a , then $\text{lcm}(a, b) = b$, so $h(b) = \text{lcm}(h(a), h(b))$, and hence $h(b)$ is a multiple of $h(a)$. Conversely, if $h(b)$ is a multiple of $h(a)$, then $h(b) = \text{lcm}(h(a), h(b))$. On the other hand, we have $h(c) = \text{lcm}(h(a), h(b))$. Since h is injective, we have $c = b$, so b is a multiple of a .

We apply the following claim for $H = h$.

Claim. Suppose $H: \mathbb{Z}_{\geq 2} \rightarrow \mathbb{Z}_{\geq 2}$ is a bijection such that a divides b if and only if $H(a)$ divides $H(b)$. Then:

1. $H(p)$ is prime if and only if p is prime;
2. $H(\prod_{i=1}^m p_i^{e_i}) = \prod_{i=1}^m H(p_i)^{e_i}$ i.e. H is completely multiplicative;
3. H preserves gcd and lcm.

Proof. We define $H(1) = 1$, and consider the bijection $H: \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$. By the conditions on H , for any $n \in \mathbb{Z}_{\geq 2}$, n and $H(n)$ have the same number of divisors. Hence $H(p)$ is prime if and only if p is prime.

Since the only prime dividing $H(p^r)$ is $H(p)$, we have $H(p^r) = H(p)^s$ for some $s \geq 1$. Counting the number of divisors, we have $s = r$, so $H(p^r) = H(p)^r$ for any prime p and $r \geq 1$.

For $a, b \in \mathbb{Z}_{>0}$, recall that $\gcd(a, b)$ is a unique positive integer satisfying the following condition: for any $c \in \mathbb{Z}_{>0}$, c divides $\gcd(a, b)$ if and only if c divides both a and b . By the condition on H , for any $c \in \mathbb{Z}_{>0}$, $H(c)$ divides $H(\gcd(a, b))$ if and only if $H(c)$ divides both $H(a)$ and $H(b)$. Hence we have $H(\gcd(a, b)) = \gcd(H(a), H(b))$.

Similarly, we have $H(\text{lcm}(a, b)) = \text{lcm}(H(a), H(b))$. Hence we have

$$\begin{aligned} H\left(\prod_{i=1}^m p_i^{e_i}\right) &= H(\text{lcm}(p_1^{e_1}, \dots, p_r^{e_r})) = \text{lcm}(H(p_1^{e_1}), \dots, H(p_r^{e_r})) = \text{lcm}(H(p_1)^{e_1}, \dots, H(p_r)^{e_r}) \\ &= \prod_{i=1}^m H(p_i)^{e_i} \end{aligned}$$

since $H(p_i)$ and $H(p_j)$ are different primes for $i \neq j$. □

Take two primes $p \neq q$, and let x, y be positive integers such that

$$g(p) + (x - 1)h(p) = g(q) + (y - 1)h(q).$$

This is possible as $h(p)$ and $h(q)$ are two distinct primes. For every $k \geq 0$, by $P(p, x + kh(q))$ and $P(q, y + kh(p))$, we have

$$\begin{cases} g(p \cdot h(x + kh(q))) = g(p) + (x + kh(q) - 1)h(p), \\ g(q \cdot h(y + kh(p))) = g(q) + (y + kh(p) - 1)h(q), \end{cases}$$

where the right hand sides are equal. By the injectivity of g , we have

$$p \cdot h(x + kh(q)) = q \cdot h(y + kh(p)).$$

So, $h(y + kh(p))$ is divisible by p for all $k \geq 0$. By the above Claim, h preserves \gcd , so

$$h(\gcd(y, h(p))) = \gcd(h(y), h(y + h(p)))$$

is divisible by p . Since $h(p)$ is a prime, y must be divisible by $h(p)$. Moreover, $h(h(p))$ is also a prime, so we have $h(h(p)) = p$. The function $h \circ h$ is completely multiplicative, so we have $h(h(n)) = n$ for every $n \geq 2$.

By $P(a, h(b))$ and $P(b, h(a))$, we have

$$\begin{cases} g(ab) = g(a \cdot h(h(b))) \stackrel{P(a, h(b))}{=} g(a) + (h(b) - 1)h(a), \\ g(ba) = g(b \cdot h(h(a))) \stackrel{P(b, h(a))}{=} g(b) + (h(a) - 1)h(b), \end{cases}$$

so

$$g(a) + h(a)(h(b) - 1) = g(ab) = g(b) + h(b)(h(a) - 1).$$

Hence $g(a) - h(a) = g(b) - h(b)$ for any $a, b \geq 2$, so $g - h$ is a constant function. By comparing the images of g and h , the difference is -1 , i.e. $g(a) - h(a) = 1$ for any $a \geq 2$.

So, we have $g(h(a)) = h(h(a)) - 1 = a - 1$. By definition,

$$f(a - 1) = h(a) = f^{g(h(a))}(1) = f^{a-1}(1).$$

By the injectivity of f , we have $f^{a-2}(1) = a - 1$ for every $a \geq 2$. From this, we can deduce inductively that $f(a) = a + 1$ for every $a \geq 1$.

